

Review and revision of Cenozoic tropical planktonic foraminiferal biostratigraphy and calibration to the Geomagnetic Polarity and Astronomical Time Scale

Bridget S. Wade^{*,1, 2}, Paul N. Pearson³, William A. Berggren^{4,5} and
Heiko Pälike⁶

*Corresponding author: Tel: +44 (0) 113 343 4945; Fax: +44 (0) 113 343 6716.

(1) M.T. Halbouty Building, Department of Geology & Geophysics, Texas A&M
University, College Station, TX 77843-3115, USA.

(2) Present address: School of Earth and Environment, University of Leeds, Woodhouse
Lane, Leeds, LS2 9JT, United Kingdom. Email: b.wade@leeds.ac.uk

(3) School of Earth and Ocean Sciences, Cardiff University, Main Building, Park Place,
Cardiff, CF10 3YE, United Kingdom. Email: pearsonp@cardiff.ac.uk

(4) Department of Geology and Geophysics, Woods Hole Oceanographic Institute,
Woods Hole, MA, 02543, USA. Email: wberggren@whoi.edu

(5) Department of Earth and Planetary Sciences, Rutgers University, New Jersey, USA.

(6) University of Southampton, National Oceanography Centre, Southampton, European
Way, Southampton SO14 3ZH, United Kingdom. Email: heiko@noc.soton.ac.uk

Abstract

Planktonic foraminifera are widely utilized for the biostratigraphy of Cretaceous and
Cenozoic marine sediments and are a fundamental component of Cenozoic
chronostratigraphy. The recent enhancements in deep sea drilling recovery, multiple

24 coring and high resolution sampling both offshore and onshore, has improved the
25 planktonic foraminiferal calibrations to magnetostratigraphy and/or modified species
26 ranges. This accumulated new information has allowed many of the planktonic
27 foraminiferal bioevents of the Cenozoic to be revised and a reassessment of the
28 planktonic foraminiferal calibrations. We incorporate these developments and
29 amendments into the existing biostratigraphic zonal scheme.

30
31 In this paper we present an amended low-latitude (tropical and subtropical) Cenozoic
32 planktonic foraminiferal zonation. We compile 187 revised calibrations of planktonic
33 foraminiferal bioevents from multiple sources for the Cenozoic and have incorporated
34 these recalibrations into a revised Cenozoic planktonic foraminiferal biochronology. We
35 review and synthesize these calibrations to both the geomagnetic polarity time scale
36 (GPTS) of the Cenozoic and astronomical time scale (ATS) of the Neogene and late
37 Paleogene. On the whole, these recalibrations are consistent with previous work;
38 however, in some cases, they have led to major adjustments to the duration of biochrons.
39 Recalibrations of the early middle Eocene first appearance datums of *Globigerinatheka*
40 *kugleri*, *Hantkenina singanoae*, *Guembelitrioides nuttalli* and *Turborotalia frontosa* have
41 resulted in large changes in the durations of Biochrons E7, E8 and E9. We have
42 introduced (upper Oligocene) Zone O7 utilizing the biostratigraphic utility of
43 ‘*Paragloborotalia*’ *pseudokugleri*. For the Neogene Period, major revisions are applied to
44 the fohsellid lineage of the middle Miocene and we have modified the criteria for
45 recognition of Zones M7, M8 and M9, with additional adjustments regarding the
46 *Globigerinatella* lineage to Zones M2 and M3. The revised and recalibrated datums

provide a major advance in biochronologic resolution and a template for future progress to the Cenozoic time scale.

Keywords: Cenozoic time scale, planktonic foraminifera, time scale calibration, Neogene time scale; Paleogene time scale, biostratigraphy

1. Introduction

1.1 Cenozoic planktonic foraminiferal biostratigraphy

Robust stratigraphic correlations are essential to decipher Earth history. Planktonic foraminifera have many characteristics considered ideal for biostratigraphic index fossils – morphologically distinct, diverse, rapidly-evolving, highly abundant, often globally distributed and high preservation potential. As such, they are extensively used for the biostratigraphy of Cretaceous and Cenozoic marine sediments and are a fundamental component of Cenozoic chronostratigraphy. Regional biostratigraphic schemes developed in parallel with taxonomic and stratigraphic research, beginning in the 1940s in the oil-producing parts of the USA (e.g. Cushman and Stainforth, 1945) and reaching high levels of sophistication by mid-century for various key economic areas in the West (e.g. Bolli 1957a, b) and Former Soviet Union (e.g. Subbotina, 1953). The process of generalizing these essentially local schemes was underway in the 1960s (e.g., Bandy, 1964; Banner and Blow, 1965) but was accelerated by the Deep Sea Drilling Project (DSDP), when it soon became clear that even in the deep Pacific Ocean there were “the same species assemblages and faunal succession as reported from other areas” (Krasheninikov, 1971, p. 1055-1056). Hence integrated biostratigraphic schemes began to be regarded as global

within broad latitudinal belts, although some degree of provincialism was also recognized in different ocean basins, the Mediterranean Sea (e.g. Cita, 1973), and areas of high productivity.

The starting point for all biostratigraphy is the recognition of so-called biostratigraphic horizons (biohorizons) that can be thought of as levels that can be correlated between stratigraphic sections where the fossil content changes in some measurable way (see McGowran, 2005 for discussion). The biostratigraphic resolution is determined by morphospecies evolution (appearances and disappearances of species). Additional biohorizons include prominent changes in coiling direction. Since at least the mid-nineteenth century (Hedberg et al., 1976) it has been standard biostratigraphic practice to divide stratigraphic sections into zones and subzones, which are non-overlapping slices of stratigraphy that lie between prominent biohorizons and are characterized by a particular fossil content (see McGowran, 2005). The zones and subzones allow recognition of fairly broad and easily identifiable intervals of stratigraphy that can be widely correlated with confidence. The practice of naming and/or sequentially numbering biozones provides the biostratigrapher with a useful mnemonic and easy means of communication. The five types of biozones that can logically be based on stratigraphic lowest and highest occurrences (LO and HO) are shown in Fig. 1.

Throughout the Cenozoic the planktonic foraminifera have been most abundant and diverse in the tropics and subtropics, hence it is for these latitudes that the zonal schemes are most detailed and easy to apply. Work in the temperate mid-latitudes and sub-polar

oceans, especially in the Neogene when climatic gradients were more pronounced than in the Paleogene, either requires judicious use of secondary markers and the amalgamation of zones when key species are absent or the development of entirely separate zonal schemes (e.g. Jenkins 1966, 1967, 1971; Poore and Berggren, 1975; Kaneps, 1975; Kennett and Srinivasan, 1983; Stott and Kennett, 1990). Nevertheless the tropical / subtropical schemes have always provided the central standard, and within that standard an increasing number of key datum levels have been accurately calibrated against magneto- and astrochronological time scales.

The updated and revised tropical / subtropical zonation presented here is a much-modified and refined descendant of that originally developed by British Petroleum micropaleontologists in the Caribbean and Tanzania in the 1950s and 1960s (Bolli, 1957a, b, 1966; Blow, 1959; Blow and Banner, 1962; Bolli and Bermúdez, 1965). A significant innovation was made by Banner and Blow (1965) who partially described a set of zones using alphanumeric shorthand ('P' for Paleogene and 'N' for Neogene, with only the Neogene zones fully described, from N1 to N22). The remainder of this scheme was presented by Blow (1969) with extensive taxonomic and stratigraphic discussion; however note that one unfortunate consequence of this was that the Neogene began with Zone N4 due to uncertainty in the placement of the Oligocene/Miocene boundary. The Blow (1969) scheme was later revised and expanded upon by Blow (1979). A variant of this zonation was published by Berggren (1969) and originally attributed to "Blow and Berggren in Berggren (1969)"; see discussion in Berggren and Miller (1988). Numerous amendments to these zones have been suggested over the years, often for specific parts of

the scheme that have failed to work optimally in certain areas. Comprehensive updates and correlations between alternative schemes were published by Stainforth et al. (1975), Kennett and Srinivasan (1983), Bolli and Saunders (1985), Berggren and Miller (1988), and Berggren et al. (1995; referred to herein as BKSA95). This latter review introduced a new development to the alphanumeric notation for epoch-level intervals, namely the introduction of ‘M’ for Miocene, ‘PL’ for Pliocene and ‘PT’ for Pleistocene in place of the earlier ‘N’ for Neogene zones. Similarly, Berggren and Pearson (2005) produced a new revision for the Paleogene zones following extensive taxonomic work on the Paleocene and Eocene planktonic foraminifera (Olsson et al., 1999; Pearson et al., 2006), extending this practice to include ‘P’ for Paleocene, ‘E’ for Eocene and ‘O’ for Oligocene.

1.2. Recent developments and the necessity for Cenozoic biostratigraphic review

In 2009 the research vessel the *JOIDES Resolution*, part of the Integrated Ocean Drilling Program (IODP) was refloated for a new campaign of ocean drilling, following a two year renovation. During IODP, as in DSDP and Ocean Drilling Program (ODP), planktonic foraminiferal biostratigraphy is extensively employed during Shipboard and post-cruise work to establish age-depth relationships.

The compilation of BKSA95 brought together the planktonic foraminifera and calcareous nannoplankton bioevents for the Cenozoic and has been frequently applied in regional and global biostratigraphy and correlations. Consistency of nomenclature is extremely sought after in biostratigraphy, and revision of zonal scheme(s) should not be embarked

upon without due consideration. However, since 1995, a number of apparent deficiencies in the tropical planktonic foraminiferal zonal schemes have been discovered through detailed biostratigraphic investigations and taxonomic developments. The enhancements in deep sea drilling recovery, multiple coring, high resolution sampling both offshore and onshore, has improved the calibrations with the magnetostratigraphy and/or modified the species ranges and allowed many of the planktonic foraminiferal bioevents to be revised. For example, detailed biostratigraphic investigations from Ceara Rise (Chaisson and Pearson, 1997; Pearson and Chaisson, 1997; Turco et al., 2002), equatorial Pacific Ocean (Wade et al., 2007), Indian Ocean (Hancock et al., 2002) and the Gorrondaxe and Agost sections in Spain (Payros et al., 2007, 2009; Ortiz et al., 2008; Larrasoña et al., 2008) have resulted in revision of the calibrations of numerous bioevents. Most of these changes are small and incremental in nature, but some (e.g., the revision of the lower middle Eocene by Payros et al., 2007) are major developments. This accumulated new information presents the opportunity for a reassessment of the planktonic foraminiferal calibrations and to incorporate developments and amendments to the existing biostratigraphic zonal scheme.

Ocean Drilling Program Leg 154 (Ceara Rise, western tropical Atlantic Ocean) produced several new constraints on tropical planktonic foraminiferal biohorizons that are incorporated into the present study. The sedimentary succession on Ceara Rise is remarkably complete and continuous from the Paleocene to Recent (Curry et al., 1995). It is unfortunate that no magnetostratigraphy is available for the sites. Nevertheless a very complete series of foraminifera and nannofossil biohorizons was recorded (Chaisson and

Pearson, 1997; Pearson and Chaisson, 1997; Turco et al., 2002; Backman and Raffi, 1997). Datums that are derived from Leg 154 sites are indirectly calibrated to the geomagnetic magnetostratigraphic time scale by interpolation between other well calibrated events and through linear interpolation with the astrochronology of Lourens et al. (2004). To ensure consistency we have also recalculated bioevents from Ceara Rise (Chaisson and Pearson, 1997; Pearson and Chaisson, 1997; Turco et al., 2002) and converted them to the magnetochronology of Cande and Kent (1995) (Table 1, Fig. 2). A revised Cenozoic planktonic foraminiferal biochronology is a logical outcome of the improved age control, and the new campaign of ocean drilling by the *JOIDES Resolution* (e.g., Pälike et al., 2009; Lyle et al., 2009), as well as high-resolution biostratigraphic studies since 1995 that have acted as the catalyst to bring these new calibrations together.

1.3. A Cenozoic Astronomical Naming Scheme

The ~405 kyr cycle of Earth's eccentricity is regarded as relatively stable over geological time (Laskar, 1999). Neogene geochronology has undergone major advances with integrated magneto-, astro-stratigraphies of continuous open marine and outcrop sedimentary successions (Hilgen et al., 2006 and references therein) and a well constrained astro-magneto-chronology for the entire Neogene (Lourens et al., 2004). An orbitally calibrated magnetochronology for the Paleogene is still under development and is available to Chron C19n (Pälike et al., 2006). We propose here a naming scheme that relates astronomical (chronological) information with magnetostratigraphy. The naming scheme to define events by ~405 kyr eccentricity cycle follows the procedure used in Wade and Pälike (2004) and Pälike et al. (2006). The cycle count number is identified by

the ~405 kyr eccentricity minima from Laskar et al. (2004), numerically coded, starting with number 1 for the most recent minimum and proceeding back in time (Table 2). As in Wade and Pälike (2004) we include a subscripted code for the geological epoch together with the magnetochron (excluding subchrons) closest to the ~405 kyr eccentricity minimum (Fig. 3).

1.4. Revision of Cenozoic tropical planktonic foraminiferal bio-, magneto-, astro-chronology

We have produced a revised and unified Cenozoic planktonic foraminiferal magnetobiochronology. We integrate planktonic foraminiferal data from multiple sources and have incorporated these recalibrations into a revised Cenozoic planktonic foraminiferal biochronology and review and synthesize these calibrations to both the geomagnetic polarity time scale (GPTS) and astronomical time scale (ATS). The biochronology has been derived from calibrations to the magnetostratigraphic polarity zones (chrons and subchrons) in deep sea and land sections where available. We have used linear interpolation to convert numerous bioevents to multiple time scales (Cande and Kent, 1995; Lourens et al., 2004; Luterbacher et al., 2004; Pälike et al., 2006) to provide the reader with convenient “look up” tables and figures for age models and biostratigraphic control. The new and former calibration ages are given in Tables 1, 3 and 4.

Here, we present an amended low-latitude (tropical and subtropical) Cenozoic planktonic foraminiferal biochronology of 187 planktonic foraminiferal events for the Cenozoic (108

Neogene and 79 Paleogene) (Figs. 2 and 4; Tables 1, 3 and 4). Bioevents have been recalibrated to the GPTS of Cande and Kent (1995) and Luterbacher et al. (2004) and to the ATS of Lourens et al. (2004) and Pälike et al. (2006, from the Oligocene/Miocene boundary to Zone E11) (Tables 1, 3 and 4). We recalibrate 61 primary and over 120 secondary bioevents for the Cenozoic. On the whole, these recalibrations are consistent with previous work. However, in some cases, they have led to major adjustments to the duration of biochrons (Figs. 2-4). Our revised and recalibrated datums provide a major advance in biochronologic resolution and a template for future progress to the Cenozoic time scale. The calibrations presented here represent the current status of Cenozoic tropical biostratigraphy and further updates and refinements are likely to follow with future IODP Expeditions. The enhanced recovery, multiple hole advanced piston coring, of successions with high sedimentation rates and paleomagnetic control (e.g., Pälike et al., 2009; Lyle et al., 2009) will enable high resolution biostratigraphic studies and extension of the ATS beyond 40 Ma. We use the PT, PL, and M zonal scheme of BKSA95 and the O, E and P zonal scheme of Berggren and Pearson (2005). The 21 zones of the Neogene Period (BKSA95) and 29 zones of the Paleogene Period (Berggren and Pearson, 2005) are retained and amended to reflect updated chronostratigraphic calibration to the GPTS.

As with previous compilations (e.g., Berggren et al., 1985, 1995b), our magnetobiochronology is founded on first order calibrations between biostratigraphic events and the magnetostratigraphy in ocean drilling cores, as well as outcrop sections. These are supplemented with orbital compilations in instances where a

magnetostratigraphy was absent (e.g., Ceara Rise). Here the current status of Cenozoic planktonic foraminiferal biostratigraphy is reviewed, refined and recalibrated, with modifications to the zonal criteria where necessary. All calibrated bioevents are listed in Tables 1, 3 and 4, primary events that define zonal boundaries are shown in bold and on Figs. 2-4.

Our revised zonation is primarily for application in open ocean settings of the Pacific, Indian and Atlantic oceans, and therefore we have not incorporated biostratigraphic information that is regionally restricted, such as to the Mediterranean, the high latitudes and other localized environments and when studies subsequent to BKSA95 have indicated them to be diachronous, unreliable or require further evaluation. Secondary bioevents that have not been used include: the last appearance datum (LAD) *Globoquadrina pseudofoliata* (Chaproniere et al., 1994), first appearance datum (FAD) *Globorotalia hirsuta* (Pujol and Duprat, 1983), FAD *Globoconella inflata* (Berggren et al., 1995a), FAD *Pulleniatina finalis* (Chaproniere et al., 1994), LAD *Neogloboquadrina atlantica* (Weaver and Clement, 1987), LAD *Globoconella puncticulata* (Atlantic) (Zijderveld et al., 1991), LAD *Globoquadrina baroemoenensis* (Curry et al., 1995), FAD *Globorotalia sphericomiozea* and *Globorotalia pliozea* (Srinivasan and Sinha, 1992), LAD *Globorotalia zealandica* (Li et al., 1992), FAD *Globigerinoides altiapertura* (Steininger et al., 1997), LAD *Globigerina labiacrassata* (BKSA95), FAD *Globigerinita boweni* (Li et al., 1992), LAD *Clavigerinella eocanica* (Pearson and Chaisson, 1997), LAD *Subbotina linaperta* (Wade, 2004), and LAD *Planorotalites capdevilensis* (Wade, 2004).

2. Neogene Period

The genus *Globorotalia* has been widely utilized in morphometric and biostratigraphic studies. Several subgenera exist in the literature for keeled forms, that have been somewhat inconsistently been applied by various workers. For example Cushman and Bermúdez (1949) named the subgenus *Globorotalia* (*Truncorotalia*) with *G. truncatulinoides* as the type species. Bandy (1972) named several other subgenera (*Menardella*, *Fohsella*, *Hirsutella*) though these were not formally described and no type species was designated. Kennett and Srinivasan (1983) used these as subgenera and designated type species. As in the Paleocene (Olsson et al., 1999) and Eocene (Pearson et al., 2006) taxonomic atlases we have chosen not to use subgenera and refer to most of the above forms as *Globorotalia*, which are all part of a single clade descended from Miocene *G. praescitula* (Kennett and Srinivasan, 1983). The exception to this is the distinct taxonomic lineage of *Fohsella* which we use at the generic level because it is very likely polyphyletic with respect to the true *Globorotalia* and has a subtly different wall texture.

For the Neogene Period, the incorporation of revised bioevents from Ceara Rise (adopted by Lourens et al., 2004) has led to some major modifications to the planktonic foraminiferal stratigraphy and zonal scheme. Lourens et al. (2004) did not discuss the implications of their revised chronology to planktonic foraminiferal biostratigraphy. We have re-evaluated and assessed these events and compared them to previous magnetostratigraphic calibrations. In the majority of events discussed below the concept

of the zone has remained the same, with the exception of Zones M9, M8, M7, M3, and M2.

Since the initiation of this work, The International Commission on Stratigraphy have proposed to lower the base of the Quaternary Period and the Pleistocene epoch to 2.58 Ma, at the same time capping the Neogene Period at that age (Gibbard et al., 2009). As this is not yet formally ratified in publication and has been met with widespread opposition (e.g., Van Couvering et al., 2009), we have used the former definition of the "real" Neogene (McGowran et al., 2009) as consisting of the Miocene, Pliocene, Pleistocene and Holocene/Recent, while awaiting a resolution of the controversy.

2.1. Amendment to the Pleistocene PT Zones

Zone PT1. *Globigerinoides ruber* Partial-range Zone (herein renamed = Zone PT1 [*Globigerinoides fistulosus*-*Globorotalia truncatulinoides* Interval Zone] of BKS95).

Definition: Biostratigraphic interval characterized by the partial range of the nominate taxon between the highest occurrence (HO) of *Globigerinoides fistulosus* and the Recent.

Magnetochronologic calibration: Chron C2n-Chron C1n (present day).

Astronomical cycle calibration: 5_{Pt}-C1_r - present day.

Estimated age: 1.88–0 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); late Pliocene-Recent.

Remarks: The definition of the zone remains the same as in BKS95, however, it has been renamed here according to the convention in Fig. 1 and of Berggren and Pearson

(2005). Lourens et al. (2004) provided two astronomical ages for the LAD of *Globigerinoides fistulosus*, 1.77 Ma from Site 677 (Shipboard Scientific Party, 1988; Shackleton et al., 1990) and 1.88 Ma (Chaisson and Pearson, 1997). The calibration from Site 677 is poorly constrained between relatively widely spaced core catcher samples and requires further investigation, thus we use the LAD of *Globigerinoides fistulosus* from Ceara Rise of 1.88 Ma.

Subzone PT1b. *Globorotalia truncatulinoides* Partial-range Subzone

Definition: Biostratigraphic interval characterized by the partial range of the nominate taxon between the highest occurrence (HO) of *Globorotalia tosaensis* and the Recent.

Magnetochronologic calibration: Chron C1n.

Astronomical cycle calibration: 2_{Pt-C1n} - present day.

Estimated age: 0.61–0 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); late Pleistocene (including Holocene).

Remarks: The nomenclature and definition of this subzone remain the same as in Berggren et al. (1995a) and BKS95.

Subzone PT1a. *Globorotalia tosaensis* Highest-occurrence Subzone (herein renamed = Subzone PT1a [*Globigerinoides fistulosus*-*Globorotalia tosaensis* Interval Sub-Zone] of Berggren et al., 1995a and BKS95).

Definition: Biostratigraphic interval between the HO of *Globigerinoides fistulosus* and the HO of the nominate taxon, *Globorotalia tosaensis*.

Magnetochronologic calibration: Chron C2n – Chron C1n.

- 323 *Astronomical cycle calibration:* 5_{Pt-C1r} - 2_{Pt-C1n}.
- 324 *Estimated age:* 1.88–0.61 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); late
- 325 Pliocene to late Pleistocene.
- 326 *Remarks:* This subzone is the same as Subzone PT1a of Berggren et al. (1995a) and
- 327 renamed according to the convention of Berggren and Pearson (2005). See discussion for
- 328 Zone PT1 regarding the LAD of *Globigerinoides fistulosus*.
- 329
- 330 *2.2. Amendment to the Pliocene PL Zones*
- 331 **Zone PL6 (Indo-Pacific). *Globigerinoides fistulosus* Highest-occurrence Zone (Indo-**
- 332 **Pacific)** (herein renamed = Zone PL6 [*Globorotalia pseudomiocenica* -
- 333 *Globigerinoides fistulosus* Interval Zone] of Berggren et al., 1995a and BKSA95).
- 334 *Definition:* Biostratigraphic interval between the HO of *Globorotalia pseudomiocenica*
- 335 and the HO of the nominate taxon, *Globigerinoides fistulosus* in the Indo-Pacific
- 336 province.
- 337 *Magnetochronologic calibration:* Subchron C2r.2r to Chron C2n.
- 338 *Astronomical cycle calibration:* 6_{Pl-C2n} - 5_{Pt-C1r}.
- 339 *Estimated age:* 2.30–1.88 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); late
- 340 Pliocene.
- 341 *Remarks:* This zone is the same as Zone PL6 (Indo-Pacific) of Berggren et al. (1995a)
- 342 and BKSA95. This zone is specific to the Indo-Pacific realm because *Globorotalia*
- 343 *pseudomiocenica* evolved into *G. miocenica* over the interval of Chron C2An.3n to
- 344 Chron C2An.2n (~ 3.5-3.2 Ma) in the Atlantic realm (DSDP Site 502, Colombia Basin;
- 345 Keigwin, 1982), whereas it persisted into younger biostratigraphic level in the Indo-

Pacific realm (BKSA95: 166). It is approximately equivalent in stratigraphic level to Zone PL6 (Atlantic). Renamed according to the convention of Berggren and Pearson (2005).

Zone PL6 (Atlantic). *Globigerinoides fistulosus* Highest-occurrence Zone (Atlantic)

(herein renamed = Zone PL6 [*Globorotalia miocenica* -*Globigerinoides fistulosus* Interval Zone] of Berggren et al., 1995a and BKSA95).

Definition: Biostratigraphic interval between the HO of *Globorotalia miocenica* and the HO of the nominate taxon, *Globigerinoides fistulosus* in the Atlantic province.

Magnetostratigraphic calibration: Subchron C2r.2r-Subchron C1r.2r.

Astronomical cycle calibration: 7_{Pl}-C2r - 5_{Pt}-C1r.

Estimated age: 2.39–1.88 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); late Pliocene.

Remarks: This zone is the same as Zone PL6 (Atlantic) of Berggren et al. (1995a) and BKSA95. This zone is specific to the Atlantic Ocean (see remarks regarding Zone PL6 [Indo-Pacific]). It is approximately equivalent in stratigraphic level to Zone PL6 (Indo-Pacific). Renamed according to the convention of Berggren and Pearson (2005).

Zone PL5 (Indo-Pacific). *Globorotalia pseudomiocenica* Highest Occurrence Zone

(herein renamed = Zone PL5 [Indo-Pacific] [*Dentoglobigerina altispira* - *Globorotalia pseudomiocenica* Interval Zone] of Berggren et al. 1995a and BKSA95).

Definition: Biostratigraphic interval between the HO of *Dentoglobigerina altispira* and

369 HO of the nominate taxon *Globorotalia pseudomiocenica* in the Indo-Pacific province.

370 *Magnetostratigraphic calibration*: Subchron C2An.3n-Subchron C2r.2r.

371 *Astronomical cycle calibration*: 9Pl-C2An - 6Pl-C2n.

372 *Estimated age*: 3.46–2.30 Ma (as per Cande and Kent, 1995); 3.47-2.30 Ma (as per

373 Lourens et al., 2004); late Pliocene.

374 *Remarks*: Renamed according to the convention of Berggren and Pearson (2005).

375

376 **Zone PL5 (Atlantic). *Globorotalia miocenica* Highest Occurrence Zone** (herein

377 renamed = Zone PL5 [Atlantic] [*Dentoglobigerina altispira* - *Globorotalia*

378 *miocenica* Interval Zone] of Berggren et al. 1995a and BKS95).

379 *Definition*: Biostratigraphic interval between the HO of *Dentoglobigerina altispira* and

380 HO of the nominate taxon *Globorotalia miocenica* in the Atlantic province.

381 *Magnetostratigraphic calibration*: Subchron C2An.2n-Subchron C2r.2r.

382 *Astronomical cycle calibration*: 8Pl-C2An - 7Pl-C2r.

383 *Estimated age*: 3.13–2.30 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); late

384 Pliocene.

385 *Remarks*: Renamed according to the convention of Berggren and Pearson (2005).

386

387 **Zone PL4. *Dentoglobigerina altispira* Highest Occurrence Zone** (herein renamed =

388 Zone PL4 [*Sphaeroidinellopsis seminulina* – *Dentoglobigerina altispira* Interval

389 Zone] of Berggren et al. 1995a and BKS95).

390 *Definition*: Biostratigraphic interval between the HO of *Sphaeroidinellopsis seminulina*

391 and HO of the nominate taxon *Dentoglobigerina altispira*.

Magnetochronologic calibration: Subchron C2An.2n (Atlantic); Subchron C2An.3n (Indo-Pacific).

Astronomical cycle calibration: 9_{Pl}-C2An - 8_{Pl}-C2An (Atlantic); 10_{Pl}-C2An - 9_{Pl}-C2An (Indo-Pacific).

Estimated age: Atlantic Ocean = 3.16–3.13 Ma (as per Cande and Kent, 1995; Lourens et al., 2004); Pacific Ocean = 3.57–3.46 Ma (as per Cande and Kent, 1995); 3.59–3.47 Ma (as per Lourens et al., 2004); late Pliocene.

Remarks: Renamed according to the convention of Berggren and Pearson (2005). A short biostratigraphic interval corresponding to 30 kyr between the LAD of *Sphaeroidinellopsis seminulina* and *Dentoglobigerina altispira* was recognized by BKSA95. Chaisson and Pearson (1997) recorded these events at approximately the same stratigraphic level at Ceara Rise, but their sampling resolution suggested a short duration as in BKSA95, hence we estimate the timing of these events at 3.13 and 3.16 consistent with the stratigraphic record at Ceara Rise (Chaisson and Pearson, 1997) and the observations of BKSA95. The duration of Biochron PL4 is estimated to be 30 kyr in the Atlantic Ocean and 110 kyr in the Pacific Ocean.

Zone PL3. *Sphaeroidinellopsis seminulina* Highest Occurrence Zone (herein renamed = Zone PL3 [*Globoquadrina altispira* - *Sphaeroidinellopsis subdehiscens* Partial Range Zone] of Berggren, 1973, [*Globorotalia margaritae* – *Sphaeroidinellopsis seminulina* Interval Zone] of Berggren et al. 1995a and BKSA95).

Definition: Biostratigraphic interval between the HO of *Globorotalia margaritae* and HO of the nominate taxon *Sphaeroidinellopsis seminulina*.

- 415 *Magnetochronologic calibration:* Chron C2Ar-Subchron C2An.2n (Atlantic); Chron
 416 C2Ar-Subchron C2An.3n (Indo-Pacific).
- 417 *Astronomical cycle calibration:* 10_{Pl}-C2An - 9_{Pl}-C2An (Atlantic); 10_{Pl}-C2An - 9_{Pl}-C2An
 418 (Indo-Pacific).
- 419 *Estimated age:* Atlantic Ocean = 3.84–3.16 Ma (as per Cande and Kent, 1995); 3.85-3.16
 420 Ma (as per Lourens et al., 2004); Pacific Ocean = 3.84-3.57 Ma (as per Cande and Kent,
 421 1995); 3.85-3.59 Ma (as per Lourens et al., 2004); early-late Pliocene.
- 422 *Remarks:* The definition of this zone is the same as that of Berggren (1973). Renamed
 423 according to the convention of Berggren and Pearson (2005).
- 424
- 425 **Zone PL2. *Globorotalia margaritae* Highest Occurrence Zone** (herein renamed = Zone
 426 PL2 [*Globorotalia margaritae* – *Sphaeroidinellopsis subdehiscens* Partial Range
 427 Zone] of Berggren, 1973, [*Globoturborotalita nepenthes* - *Globorotalia*
 428 *margaritae* Interval Zone] of Berggren et al. 1995a and BKS95).
- 429 *Definition:* Biostratigraphic interval between the HO of *Globoturborotalita nepenthes*
 430 and HO of the nominate taxon *Globorotalia margaritae*.
- 431 *Magnetochronologic calibration:* Subchron C3n.1r-Chron C2Ar.
- 432 *Astronomical cycle calibration:* 12_{Pl}-C3n - 10_{Pl}-C2An.
- 433 *Estimated age:* 4.36–3.84 Ma (as per Cande and Kent, 1995); 4.37-3.85 Ma (as per
 434 Lourens et al., 2004); early Pliocene.
- 435 *Remarks:* Renamed according to the convention of Berggren and Pearson (2005).
- 436
- 437 **Zone PL1. *Globorotalia tumida* / *Globoturborotalita nepenthes* Concurrent-range**

438 **Zone** (herein renamed = Zone PL1 [*Globigerina nepenthes* - *Globorotalia tumida*
 439 Partial Range Zone] of Berggren, 1973, [*Globorotalia tumida* –
 440 *Globoturborotalita nepenthes* Interval Zone] of Berggren et al. 1995a and
 441 BKSA95).

442 *Definition:* Biostratigraphic interval between the lowest occurrence (LO) of *Globorotalia*
 443 *tumida* and HO of *Globoturborotalita nepenthes*.

444 *Magnetochronologic calibration:* Chron C3r-Subchron C3n.1r.

445 *Astronomical cycle calibration:* 15_{Mi}-C3r - 12_{Pl}-C3n.

446 *Estimated age:* Atlantic Ocean = 5.63–4.36 Ma (as per Cande and Kent, 1995); 5.72-4.37
 447 Ma (as per Lourens et al., 2004); Pacific Ocean = 5.51-4.36 Ma (as per Cande and Kent,
 448 1995); 5.57-4.37 Ma (as per Lourens et al., 2004); late Miocene-early Pliocene.

449 *Remarks:* Renamed according to the convention of Berggren and Pearson (2005). The
 450 extinction of *Globorotalia cibaoensis* was used to subdivide Zone PL1 and had a
 451 calibration of 4.6 Ma in BKSA95. However, Chaisson and Pearson (1997) reported a
 452 much younger LAD for this species which was adopted by Lourens et al. (2004) to give
 453 an astronomical age on 3.23 Ma. As the much younger LAD at Ceara Rise is yet to be
 454 confirmed we use the 4.6 Ma calibration of BKSA95. Due to this discrepancy, we have
 455 removed the subdivision of Zone PL1, pending further investigations.

456

457 Consistent with previous studies (e.g., Berggren, 1977; Srinivasan and Kennett, 1981b;
 458 Kennett and Srinivasan, 1983), there is a short stratigraphic interval (32 kyr) between the
 459 HO of *Globoquadrina dehiscens* and the LO of *Sphaeroidinella dehiscens* (see Fig. 2a,
 460 Table 1). The LO of *Globorotalia tumida* occurs between these two distinctive events

(Srinivasan and Chaturvedi, 1992). The FAD *Globorotalia tumida* has been revised to 5.63 and 5.51 Ma for the Atlantic and Pacific oceans respectively (Table 1). Thunell (1981), Srinivasan and Kennett (1981a) and Chaisson and Leckie (1993) record the FAD of *G. tumida* to be older than the LAD *G. dehiscens*. However, Hodell and Kennett (1986) have shown the LAD of *G. dehiscens* to be diachronous, and the extinction appears to occur earlier in higher latitudes in comparison to tropical sites (Srinivasan and Kennett, 1981b).

2.3. Amendment to the Miocene M Zones

It is remarkable how few low latitude open ocean sections exist with good recovery, high sedimentation rates, abundant planktonic foraminifera and a clearly defined magnetostratigraphy through the Miocene. This has significantly hindered direct correlations to the GPTS and the development of robust planktonic foraminifera magnetobiostratigraphy. Miller et al. (1985) produced a magnetobiostratigraphy DSDP Sites 563 and 558 (western North Atlantic Ocean), however even these records have unconformities. Many of the events through the Miocene have been calibrated from the Buff Bay, Jamaica (BKSA95) and not from deep sea cores. Following Lourens et al. (2004) we have recalibrated the ages from Ceara Rise, but note that these are not tied to a magnetostratigraphy, and the resulting ages are significantly younger than those recorded in BKSA95. Our recalibrations have led to major changes to the age assignments and duration of Biochrons M10 to M13. We highlight some of the major changes below, but emphasize that this interval requires detailed study to confirm the ages of the events as defined from Ceara Rise (Chaisson and Pearson, 1997; Turco et al., 2002). The lower-

middle Miocene interval is divided on the diagnostic index genus *Praeorbulina*.

However, these taxa were rare at Ceara Rise (Pearson and Chaisson, 1997) and therefore were not included in the revised calibration, and we have retained the ages reported in BKSA95 for FAD of *Orbulina suturalis* (15.1 Ma), FAD *Praeorbulina circularis* (16.0 Ma), FAD *Praeorbulina curva* (16.3 Ma) and *Praeorbulina sicana* (16.4 Ma).

Zone M14. *Globigerinoides extremus* Partial-range Zone (herein renamed = Zone M14 [*Globorotalia linguaensis* - *Globorotalia tumida* Interval Zone] of BKSA95).

Definition: Partial range of the nominate taxon between the HO of *Globorotalia linguaensis* and LO of *Globorotalia tumida*.

Magnetochronologic calibration: Subchron C3An.1n-Chron C3r.

Astronomical cycle calibration: 16_{Mi}-C3r - 15_{Mi}-C3r.

Estimated age: Atlantic Ocean = 6.00–5.63 Ma (as per Cande and Kent, 1995); 6.13-5.72 Ma (as per Lourens et al., 2004); Pacific Ocean = 6.00-5.51 Ma (as per Cande and Kent, 1995); 6.13-5.57 Ma (as per Lourens et al., 2004); late Miocene.

Remarks: Following the detailed biostratigraphic investigations by Turco et al. (2002), Lourens et al. (2004) significantly revised the LAD of *Globorotalia linguaensis* to 8.97 Ma. The revised age is appreciably older than reported in BKSA95 (derived from the Tonga Plateau; Chaproniere et al., 1994) and would place the event within the *Neogloboquadrina acostaensis* Lowest-occurrence Subzone (Subzone M13a), inconsistent with the established order of bioevents. However, it should be noted that Zhang et al. (1993) found the HO of *G. linguaensis* near the same horizon as the HO *G. plesiotumida* (8.52 Ma; Table 1), which is more consistent with the older age suggested

by Turco et al. (2002). For stability we have retained the age established in BKSA95, but this interval clearly requires further investigation. The zone is renamed according to the convention of Berggren and Pearson (2005).

Zone M13. *Neogloboquadrina acostaensis* / *Globorotalia linguaensis* Concurrent-range Zone

Definition: Biostratigraphic interval between the LO of *Neogloboquadrina acostaensis* and HO of *Globorotalia linguaensis*.

Magnetostratigraphic calibration: Subchron C5n.1n-Subchron C3An.1n.

Astronomical cycle calibration: 25_{Mi}-C4Ar - 16_{Mi}-C3r.

Estimated age: 9.79–6.00 Ma (as per Cande and Kent, 1995); 9.83-6.13 Ma (as per Lourens et al., 2004); late Miocene.

Remarks: The nomenclature and definition of the zone remain the same as in BKSA95.

Subzone M13b. *Globorotalia plesiotumida* / *Globorotalia linguaensis* Concurrent-range Subzone (herein amended and renamed, approximately equivalent to Subzone M13b [*Globigerinoides extremus* / *Globorotalia plesiotumida* – *Globorotalia linguaensis* Interval Subzone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Globorotalia plesiotumida* and HO of *Globorotalia linguaensis*.

Magnetostratigraphic calibration: Subchron C4r.2r-Subchron C3An.1n.

Astronomical cycle calibration: 22_{Mi}-C4r - 16_{Mi}-C3r.

Estimated age: 8.52–6.00 Ma (as per Cande and Kent, 1995); 8.58-6.13 Ma (as per

530 Lourens et al., 2004); late Miocene.

531 *Remarks:* The definition used here removes the operational ambiguity inherent in the
 532 “and/or” designation in BKSA95, in which the Subzone was defined as the
 533 biostratigraphic interval between the LO of *Globigerinoides extremus* and/or the LO of
 534 *Globorotalia plesiotumida* and the HO of *Globorotalia linguaensis*. Evidence since
 535 BKSA95 indicates that the LOs of *Globigerinoides extremus* and *Globorotalia*
 536 *plesiotumida* may occur at different levels (compare Chaisson and Pearson, 1997 and
 537 Turco et al. 2002). The subzone is renamed here according to the convention of Berggren
 538 and Pearson (2005).

539

540 **Subzone M13a. *Neogloboquadrina acostaensis* Lowest-occurrence Subzone** (herein
 541 amended and renamed, approximately equivalent to Subzone M13a
 542 [*Neogloboquadrina acostaensis* - *Globigerinoides extremus* / *Globorotalia*
 543 *plesiotumida* Interval Subzone] of BKSA95).

544 *Definition:* Biostratigraphic interval between the LO of the nominate taxon
 545 *Neogloboquadrina acostaensis* and LO of *Globorotalia plesiotumida*.

546 *Magnetochronologic calibration:* Subchron C5n.1n-Subchron C4r.2r.

547 *Astronomical cycle calibration:* 25_{Mi}-C4Ar - 22_{Mi}-C4r.

548 *Estimated age:* 9.79–8.52 Ma (as per Cande and Kent, 1995); 9.83-8.58 Ma (as per
 549 Lourens et al., 2004); late Miocene.

550 *Remarks:* The definition used here removes the operational ambiguity inherent in the
 551 “and/or” designation in BKSA95, in which the Subzone was defined as the
 552 biostratigraphic interval between the LO of *Neogloboquadrina acostaensis* and the LO of

Globigerinoides extremus and/or the LO of *Globorotalia plesiotumida*. See remarks for Subzone M13b regarding the LADs of *Globigerinoides extremus* and *Globorotalia plesiotumida*. The subzone is renamed here according to the convention of Berggren and Pearson (2005).

There is a significant reduction in the duration of Sub-biochron M13a. The cyclostratigraphic age of the LO of *Neogloboquadrina acostaensis* (9.83 Ma) is derived from Ceara Rise (Chaisson and Pearson, 1997). This calibration was adopted by Lourens et al. (2004) and is significantly younger (1.07 myr) than in BKSA95 (10.90 Ma) and moves this event from early Subchron C5n.2n to Subchron C5n.1n. In BKSA95 the duration of this sub-biochron is 2.6 myr, from 10.9 to 8.3 Ma. Following Chaisson and Pearson (1997) this sub-biochron is recalibrated to 9.79 to 8.52 Ma and results in a change in the duration of Sub-biochron M13a to 1.27 myr, a reduction of 1.3 myr. Turco et al. (2002) noted the diachrony of the LO of *Neogloboquadrina acostaensis* between low latitudes and the Mediterranean. The age used in BKSA95 is calibrated to the magnetostratigraphy at Site 563 (Miller et al., 1985) and the discrepancy in calibrated ages may be due to further diachrony between the tropical and subtropical Atlantic Ocean, however, we note that the order of bioevents is consistent between Ceara Rise and Site 563.

Zone M12. *Globigerinoides trilobus* Partial-range Zone (herein renamed = Zone M12 [*Neogloboquadrina mayeri* – *Neogloboquadrina acostaensis* Interval Zone] of BKSA95).

576 *Definition:* Partial range of the nominate taxon between the HO of *Paragloborotalia*
 577 *mayeri* and LO of *Neogloboquadrina acostaensis*.
 578 *Magnetochronologic calibration:* Subchron C5n.2n-Subchron C5n.1n.
 579 *Astronomical cycle calibration:* $^{27}\text{Mi-C5n}$ - $^{25}\text{Mi-C4Ar}$.
 580 *Estimated age:* 10.53–9.79 Ma (as per Cande and Kent, 1995); 10.46-9.83 Ma (as per
 581 Lourens et al., 2004); late Miocene.
 582 *Remarks:* Renamed according to the convention of Berggren and Pearson (2005). As for
 583 *Neogloboquadrina acostaensis* (discussed above), there is a large difference between the
 584 age established in BKSA95 and that at Ceara Rise for the LAD of *Paragloborotalia*
 585 *mayeri* (10.53 Ma, this study; 11.40 Ma, BKSA95). The extinction of *Paragloborotalia*
 586 *mayeri* has been recalibrated to 10.53 Ma as per Chaisson and Pearson (1997) (given as
 587 *siakensis* in Turco et al., 2002). This is significantly younger (870 kyr) than the previous
 588 reported age of 11.40 Ma in BKSA95. The interpolated age would place this event mid
 589 C5n.2n rather than C5r.2r. The age used in BKSA95 is calibrated to the
 590 magnetostratigraphy at Site 563 (Miller et al., 1985) and this discrepancy may be due to
 591 diachrony between the tropical and subtropical Atlantic Ocean. Hilgen et al. (2000) noted
 592 the diachrony in the extinction of *P. mayeri* between the tropical Atlantic Ocean and the
 593 Mediterranean and diachrony with higher latitudes was suggested by Miller et al. (1991).
 594
 595 The extinction of *Paragloborotalia mayeri/siakensis* and the LO of *Neogloboquadrina*
 596 *acostaensis* and *Fohsella peripheroronda*, have been shown to be diachronous between
 597 the Mediterranean and equatorial Atlantic Ocean (Turco et al., 2002). We suggest that the
 598 younger calibrations through this interval may be due to further diachronism of extinction

events in the Jamaican sections. Clearly, further work is required to constrain the bioevents through this interval.

Zone M11. *Globoturborotalita nepenthes* / *Paragloborotalia mayeri* Concurrent-range

Zone (herein renamed = Zone M11 [*Globoturborotalita*

nepenthes/*Neogloboquadrina mayeri* Concurrent Range Zone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Globoturborotalita nepenthes* and the HO of *Paragloborotalia mayeri*.

Magnetochronologic calibration: Subchron C5r.3r-Subchron C5n.2n.

Astronomical cycle calibration: $^{29}\text{Mi-C5r}$ - $^{27}\text{Mi-C5n}$.

Estimated age: 11.55–10.53 Ma (as per Cande and Kent, 1995); 11.63-10.46 Ma (as per Lourens et al., 2004); middle-late Miocene.

Remarks: Renamed to reflect inclusion of the species *mayeri* in the genus

Paragloborotalia. In Table 1, we have incorporated the age established for the LAD of

Cassigerinella chipolensis by Turco et al. (2002). This is younger than the suggested age

of this event as in Chaisson and Leckie (1993) but appears to be a useful secondary event

within the *Globoturborotalita nepenthes* / *Paragloborotalia mayeri* Concurrent-range

Zone (Zone M11). The HO of *Globigerinoides subquadratus* is found to be near-

synchronous between Site 926 (equatorial Atlantic Ocean, Turco et al., 2002) and the

Mediterranean (Hilgen et al., 2000).

Zone M10. *Globigerinella praesiphonifera* Partial-range Zone (herein amended and

renamed = Zone M10 [*Globorotalia robusta* – *Globoturborotalita nepenthes*

Interval Zone] of BKSA95).

Definition: Partial range of the nominate taxon between the HO of *Fohsella fohsi* and LO of *Globoturborotalita nepenthes*.

Magnetochronologic calibration: Subchron C5r.3r.

Astronomical cycle calibration: $^{30}\text{Mi-C5r}$ - $^{29}\text{Mi-C5r}$.

Estimated age: 11.71–11.55 Ma (as per Cande and Kent, 1995); 11.79–11.63 Ma (as per Lourens et al., 2004); middle Miocene.

Remarks: Amended to reflect the use of the LO of *Fohsella fohsi* rather than the LO of *Fohsella lobata* and *F. robusta* as in BKSA95. Renamed according to the convention of Berggren and Pearson (2005).

Following Turco et al. (2002), Lourens et al. (2004) revised the LAD *Globorotalia praescitula* from 11.9 (BKSA95) to 13.73 Ma. This placed the extinction of *G. praescitula* between the LOs of *F. 'praefohsi'* and *F. fohsi (s.l.)* and thus moved the extinction of *G. praescitula* from Zone M10 to M7. Initial investigations from Site U1337 (Shipboard Scientific Party, in press) suggest that the biostratigraphic events are consistent with BKSA95 and therefore we have retained the calibration of 11.9 Ma here pending further investigations.

2.3.1. Revision of Zones M7-M9 (*Fohsella lineage*)

One of the key lineages used in the biostratigraphic subdivision of the middle Miocene is the fohsellid lineage (usually referred to in the earlier literature as *Globorotalia fohsi* and its various subspecies). All previous work agrees that early representatives of the lineage

tend to be small forms with rounded peripheries, and that there is a gradual trend through time towards larger size and more acute peripheries which eventually results in keeled forms. After this more lobate morphotypes appear, as do more robust, biconvex forms. Taxonomic subdivision of this gradual chronocline into species and subspecies is inevitably subjective, and contrasting approaches were taken by Bolli (1957b), Blow (1957), Blow and Banner (1966), Olsson (1972), Stainforth et al. (1975), Kennett and Srinivasan (1983), Bolli and Saunders (1985) and Berggren (1993). These various taxonomic schemes are necessarily mirrored in different approaches to the biostratigraphic subdivision of the middle Miocene based upon the taxa.

The biostratigraphic scheme used here (in slightly modified form; see discussion below) is that first suggested by Banner and Blow (1965), Blow and Banner (1966) and N zonal concepts of Blow (1969). This scheme capitalizes on the biostratigraphic utility of this evolutionary lineage and uses the successive first occurrences of *F. peripheroacuta* (a morphotype that has a distinctly pinched or acute periphery), *F. 'praefohsi'* (which has a incipient keel on the final one or two chambers), and *F. fohsi* (which has a well-developed keel on the final chamber and an incipient keel throughout the last whorl), and finally the extinction of the group as successive zonal boundaries. These biohorizons originally delimited Zones N10-N12 (see also Kennett and Srinivasan, 1983) and in our scheme they delimit Zones M7-M9 (Fig. 5). We have found this scheme to provide excellent biostratigraphic control in the tropical Pacific and Atlantic Oceans (Pearson, 1995; Pearson and Chaisson, 1997).

There is, however, a taxonomic problem, as highlighted by Bolli and Saunders (1985), in that the morphology of the holotype of *Globorotalia (Fohsella) praefohsi* Blow and Banner does not seem to accord well with the *concept* of the taxon as originally suggested by Blow and Banner (1966). We agree with Bolli and Saunders that the *praefohsi* holotype can be regarded as a subjective synonym of *Globorotalia lobata* Bermúdez (= *Fohsella lobata*), which is a more ‘advanced’ member of the lineage. The paratype, however, is more in accord with the concept of *praefohsi* as originally articulated by Blow and Banner (1966). It is also pertinent that the holotype of *praefohsi* comes from a higher stratigraphic level than the paratype (Bolli and Saunders, 1985). Pearson (1995) and Pearson and Chaisson (1997) acknowledged this problem by referring to the N11 Zone fossil informally as *Fohsella ‘praefohsi’*, as we have done, thereby retaining the concept of Blow and Banner (1966) as widely used subsequently (e.g. Kennett and Srinivasan, 1983) and deferring resolution of the taxonomic problem for future work. In contrast, BKSA95 placed *praefohsi* in synonymy with *lobata* and effectively eliminated the old biostratigraphic subdivision between Zones N11 and N12. At the same time BKSA95 introduced the first alphanumeric M-zone scheme for the Miocene and used the first occurrence of the most ‘advanced’ of all the fohsellids, *F. robusta* as a subzone marker for the first time (delimiting their Subzones M9a and M9b).

The problem with the solution of BKSA95 is that it eliminates a proven highly useful zone fossil (*F. ‘praefohsi’*) along with the biostratigraphic resolution that it provides. It is clear that detailed taxonomic revision of *praefohsi* is required, based on new SEM micrographs of the relevant types and detailed descriptions. If the holotype does indeed

prove to be a synonym of *lobata*, as the illustration suggests, then it may be desirable to name a new species that accords better with the intended concept of *praefohsi*, which could then be the zone fossil for Zone M8. It may be that Blow and Banner's paratype could be used to typify such a new species, or alternatively a holotype could be taken from well-preserved and well-dated assemblages from elsewhere. Pending such a resolution, we retain the informal taxon *F. 'praefohsi'* as the zone fossil for Zone M8 basing our concept not on the holotype illustration but on Blow and Banner's original description as well as subsequent illustrated specimens that accord with that concept such as those shown by Kennett and Srinivasan (1983).

Zone M9. *Fohsella fohsi* Taxon-range Zone (herein defined, approximately equivalent to Zone N12 [*Globorotalia (G.) fohsi* Partial-range Zone] of Blow, 1969; combined Zone M9 [*Globorotalia fohsi lobata* - *Globorotalia fohsi robusta* Interval Zone] and Zone M8 [*Globorotalia fohsi sensu stricto* Lineage Zone] of BKS95).

Definition: Biostratigraphic interval characterized by the total range of the nominate taxon between its LO and HO.

Magnetostratigraphic calibration: Chron C5ABn-Subchron C5r.3r.

Astronomical cycle calibration: $^{34}\text{Mi-C5AAr}$ - $^{30}\text{Mi-C5r}$.

Estimated age: 13.34–11.71 Ma (as per Cande and Kent, 1995); 13.41–11.79 Ma (as per Lourens et al., 2004); middle Miocene.

Remarks: The new concept of Zone M9 follows from our modification of middle Miocene zonation discussed above. The HO of *Fohsella robusta* and *F. fohsi* are

estimated to be at the same stratigraphic level. However, in the astronomical calibrations based on Ceara Rise, Turco et al. (2002) did not differentiate species within the fohsellid group and Chaisson and Pearson (1997) did not comment on the HO of *F. robusta*. Therefore we use HO of *F. fohsi* to define the top of Zone M9.

Subzone M9b. *Fohsella robusta* / *Fohsella fohsi* Concurrent-range Subzone (herein amended and renamed = Subzone M9b [*Globorotalia fohsi robusta* Total Range Zone] of BKSA95; [*Globorotalia fohsi robusta* Total Range Zone] of Bolli, 1957b).

Definition: Biostratigraphic interval between the LO of *Fohsella robusta* and the HO of *Fohsella fohsi*.

Magnetochronologic calibration: Chron C5AAn-Subchron C5r.3r.

Astronomical cycle calibration: 33_{Mi}-C5Ar - 30_{Mi}-C5r.

Estimated age: 13.09–11.71 Ma (as per Cande and Kent, 1995); 13.13-11.79 Ma (as per Lourens et al., 2004); middle Miocene.

Remarks: The concept of Subzone M9b follows from our modification of middle Miocene zonation discussed above.

Subzone M9a. *Fohsella fohsi* Lowest-occurrence Subzone (herein defined, equivalent to Zone M8 [*Globorotalia fohsi sensu stricto* Lineage Zone] and Subzone M9a [*Globorotalia fohsi lobata* Lineage Zone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Fohsella fohsi* and the LO of *Fohsella robusta*.

737 *Magnetostratigraphic calibration:* Chron C5ABn-Chron C5AAn.

738 *Astronomical cycle calibration:* $^{34}\text{Mi-C5AAr}$ - $^{33}\text{Mi-C5Ar}$.

739 *Estimated age:* 13.34-13.09 Ma (as per Cande and Kent, 1995); 13.41-13.13 Ma (as per

740 Lourens et al., 2004); middle Miocene.

741 *Remarks:* The new concept of Subzone M9a follows from our modification of middle

742 Miocene zonation discussed above. The subzone differs from the concept of Subzone

743 M9a (*Globorotalia fohsi lobata* Lineage Zone) as per BKS95, which was defined as the

744 biostratigraphic interval between LO of *Fohsella lobata* and LO of *Fohsella robusta*. We

745 have been unable to utilize the LO of *F. lobata* as Chaisson and Pearson (1997) and

746 Turco et al. (2002) did not provide stratigraphic constraints on this taxon from Ceara Rise

747 and therefore there is presently no astronomical calibration (Lourens et al., 2004). This is

748 unfortunate as *F. lobata* is a distinctive and the LO of this taxon appears isochronous in

749 low latitudes (Srinivasan and Chaturvedi, 1992).

750

751 Turco et al. (2002) provided a refined calibration for the LAD of *Cassiginella*

752 *martinezpicoi* at Ceara Rise, which is consistent with studies elsewhere (Chaisson and

753 Leckie, 1993). We have incorporated the revised age for the LAD of *Cassiginella*

754 *martinezpicoi* in tables 1 and 3.

755

756 **Zone M8. *Fohsella* ‘*praefohsi*’ Lowest-occurrence Zone** (herein defined = Zone N11

757 [*Globorotalia* (*G.*) *praefohsi* Consecutive-range Zone] of Blow, 1969; and the

758 upper part of Zone M7 [*Globorotalia peripheroacuta* Lineage Zone] of

759 BKS95).

760 *Definition:* Biostratigraphic interval between the LO of the nominate taxon *Fohsella*
761 *'praefohsi'* and the LO of *Fohsella fohsi*.
762 *Magnetostratigraphic calibration:* Chron C5ACn-Chron C5ABn.
763 *Astronomical cycle calibration:* $^{35}\text{Mi-C5ABr}$ - $^{34}\text{Mi-C5AAr}$.
764 *Estimated age:* 13.74–13.34 Ma (as per Cande and Kent, 1995); 13.77-13.41 Ma (as per
765 Lourens et al., 2004); middle Miocene.
766 *Remarks:* The new concept of Zone M8 follows from our modification of middle
767 Miocene zonation discussed above.
768
769 **Zone M7. *Fohsella peripheroacuta* Lowest-occurrence Zone** (herein defined = Zone
770 N10 [*Globorotalia (Turborotalia) peripheroacuta* Consecutive-range Zone] of
771 Blow, 1969; and the lower part of Zone M7 [*Globorotalia peripheroacuta*
772 Lineage Zone] of BKSA95).
773 *Definition:* Biostratigraphic interval between the LO of the nominate taxon *Fohsella*
774 *peripheroacuta* and the LO of *Fohsella 'praefohsi'*.
775 *Magnetostratigraphic calibration:* Chron C5ADn-Chron C5ACn.
776 *Astronomical cycle calibration:* $^{36}\text{Mi-C5ACn}$ - $^{35}\text{Mi-C5ABr}$.
777 *Estimated age:* 14.23–13.74 Ma (as per Cande and Kent, 1995); 14.24-13.77 Ma (as per
778 Lourens et al., 2004); middle Miocene.
779 *Remarks:* The new concept of Zone M7 follows from our modification of middle
780 Miocene zonation discussed above. Turco et al. (2002) provided a refined calibration for
781 the LAD *Globorotalia archeomenardii* at Ceara Rise, which is consistent with studies

elsewhere (Chaisson and Leckie, 1993). We have incorporated the refined calibration for the LAD *Globorotalia archeomenardii* in tables 1 and 3.

Zone M6. *Orbulina suturalis* Lowest-occurrence Zone (herein renamed = Zone M6 [*Globorotalia peripheroronda* Partial-range Zone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Orbulina suturalis* and the LO of *Fohsella peripheroacuta*.

Magnetochronologic calibration: Subchron C5Bn.2n-Chron C5ADn.

Astronomical cycle calibration: $^{38}\text{Mi-C5Bn}$ - $^{36}\text{Mi-C5ACn}$.

Estimated age: 15.10-14.23 Ma (as per Cande and Kent, 1995); 15.10-14.24 Ma (as per Lourens et al., 2004); middle Miocene.

Remarks: Because of the rarity of *Orbulina* at the beginning of its range at Ceara Rise (Pearson and Chaisson, 1997) we have retained the age estimate from BKSA95. The Zone is renamed according to the convention of Berggren and Pearson (2005).

Zone M5. *Praeorbulina sicana* Lowest-occurrence Zone (herein renamed = Zone M5 [*Praeorbulina sicana* – *Orbulina suturalis* Interval Zone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Praeorbulina sicana* and the LO of *Orbulina suturalis*.

Magnetochronologic calibration: Subchron C5Cn.2n-Subchron C5Bn.2n.

Astronomical cycle calibration: $^{41}\text{Mi-C5Cn}$ - $^{38}\text{Mi-C5Bn}$.

Estimated age: 16.40-15.10 Ma (as per Cande and Kent, 1995); 16.38-15.10 Ma (as per Lourens et al., 2004); early-middle Miocene.

Remarks: Renamed according to the convention of Berggren and Pearson (2005). We follow the criterion of Jenkins et al. (1981) for the identification of *sicana*, see also discussion in Kennett and Srinivasan (1983) and Pearson (1995).

Subzone M5b. *Praeorbulina glomerosa* Lowest-occurrence Subzone (herein renamed = Subzone M5b [*Praeorbulina glomerosa sensu stricto* – *Orbulina suturalis* Interval Subzone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Praeorbulina glomerosa* and the LO of *Orbulina suturalis*.

Magnetostratigraphic calibration: Subchron C5Cn.1r-Subchron C5Bn.2n.

Astronomical cycle calibration: 41_{Mi}-C5Cn - 38_{Mi}-C5Bn.

Estimated age: 16.29-15.10 Ma (as per Cande and Kent, 1995); 16.27-15.10 Ma (as per Lourens et al., 2004); early-middle Miocene.

Remarks: Renamed according to the convention of Berggren and Pearson (2005).

Subzone M5a. *Praeorbulina sicana* Lowest-occurrence Subzone (herein renamed = Subzone M5a [*Praeorbulina sicana* - *Praeorbulina glomerosa sensu stricto* Interval Subzone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Praeorbulina sicana* and the LO of *Praeorbulina glomerosa*.

Magnetostratigraphic calibration: Subchron C5Cn.2n-Subchron C5Cn.1r.

Astronomical cycle calibration: 41_{Mi}-C5Cn.

Estimated age: 16.40-16.29 Ma (as per Cande and Kent, 1995); 16.38-16.27 Ma (as per

828 Lourens et al., 2004); early Miocene.

829 *Remarks:* Renamed according to the convention of Berggren and Pearson (2005).

830

831 **Zone M4. *Globigerinoides bisphericus* Partial-range Zone** (herein renamed = Zone M4

832 [*Catapsydrax dissimilis* - *Praeorbulina sicana* Interval Subzone] of BKSA95).

833 *Definition:* Partial range of the nominate taxon between the HO of *Catapsydrax dissimilis*

834 and the LO of *Praeorbulina sicana*.

835 *Magnetostratigraphic calibration:* Chron C5Dr-Subchron C5Cn.2n.

836 *Astronomical cycle calibration:* $44_{\text{Mi-C5Dn}}$ - $41_{\text{Mi-C5Cn}}$.

837 *Estimated age:* 17.62-16.40 Ma (as per Cande and Kent, 1995); 17.54-16.38 Ma (as per

838 Lourens et al., 2004); early Miocene.

839 *Remarks:* Renamed according to the convention of Berggren and Pearson (2005).

840

841 **Subzone M4b. *Fohsella birnageae* Lowest Occurrence Subzone** (herein renamed =

842 Subzone M4b [*Globigerinoides bisphericus* Partial-range Subzone] of BKSA95).

843 *Definition:* Biostratigraphic interval between the LO of *Fohsella birnageae* and the LO of

844 *Praeorbulina sicana*.

845 *Magnetostratigraphic calibration:* Subchron C5Cn.3n-Subchron C5Cn.2n.

846 *Astronomical cycle calibration:* $42_{\text{Mi-C5Cn}}$ - $41_{\text{Mi-C5Cn}}$.

847 *Estimated age:* 16.70-16.40 Ma (as per Cande and Kent, 1995); 16.69-16.38 Ma (as per

848 Lourens et al., 2004); early Miocene.

849 *Remarks:* Renamed according to the convention of Berggren and Pearson (2005).

850

851 **Subzone M4a. *Dentoglobigerina venezuelana* Partial-range Subzone** (herein renamed
852 = Subzone M4a [*Catapsydrax dissimilis* – *Globorotalia birnageae* Interval
853 Subzone] of BKSA95).

854 *Definition:* Partial range of the nominate taxon between the HO of *Catapsydrax dissimilis*
855 and the LO of *Globorotalia birnageae*.

856 *Magnetochronologic calibration:* Chron C5Dr-Subchron C5Cn.3n.

857 *Astronomical cycle calibration:* 44_{Mi}-C5Dn - 42_{Mi}-C5Cn.

858 *Estimated age:* 17.62-16.70 Ma (as per Cande and Kent, 1995); 17.54-16.69 Ma (as per
859 Lourens et al., 2004); early Miocene.

860 *Remarks:* Srinivasan and Chaturvedi (1992) point out the usefulness of the HO of
861 *Catapsydrax dissimilis* as a dissolution resistant form recorded in sites from the tropics to
862 the subantarctic. The HO of *C. dissimilis* is interpolated to within Chron C5Dr. This is
863 consistent with studies at Site 608 (Miller et al., 1991), but inconsistent with Sites 516
864 and 558 where the HO of this species is reported within C5Dn (Berggren et al., 1983).
865 Further work is required to confirm the age derived from Ceara Rise (Shackleton et al.,
866 1999). Renamed according to the convention of Berggren and Pearson (2005).

867

868 2.3.2. *Globigerinatella* spp. and *G. insueta*

869 Cushman and Stainforth (1945) initially described the genus *Globigerinatella* from the
870 Cipero Formation of Trinidad, with *G. insueta* as its only species, and used its first
871 occurrence as the marker for the base of their *Globigerinatella insueta* Zone. The first
872 occurrence of *G. insueta* remained a key zonal boundary through a number of

subsequent, more highly subdivided biostratigraphic schemes (e.g. Bolli, 1957b; Banner and Blow, 1965; Blow 1969; Kennett and Srinivasan, 1984; Bolli and Saunders, 1985).

Chaisson and Leckie (1993) were the first to describe distinct evolutionary trends in *Globigerinatella* based on their observations at ODP Site 806 on the Ontong Java Plateau, western tropical Pacific Ocean. They observed that the number of areal apertures tends to increase up section and that in the more advanced forms the apertures tend to be localized in patches on the test. Pearson (1995) described and illustrated similar evolutionary trends in specimens from ODP Sites 871 and 873 in the Marshall Islands region of the western tropical Pacific Ocean. He also observed that the earliest representatives of *Globigerinatella* all lack supplementary apertures, an observation that was confirmed by Pearson and Chaisson (1997) from ODP Sites 925 and 926 on the Ceara Rise, western tropical Atlantic Ocean.

Pearson (1995) suggested that evolutionary trends in *Globigerinatella* might be useful for the biostratigraphic subdivision of the lower Miocene, an interval which is otherwise problematic for planktonic foraminifer biostratigraphy. He suggested splitting *Globigerinatella* into two taxa based on the presence or absence of areal apertures. From observations made on Cushman and Stainforth's (1945) type material at the US National Museum, Pearson (1995) reported that the holotype and all paratypes of *G. insueta* possess areal apertures, even though they were not mentioned in the original description and are not visible on all the type illustrations. Hence forms with areal apertures were

included by Pearson (1995) and subsequently by Pearson and Chaisson (1997) in *G. insueta sensu stricto* and forms without were included as *Globigerinatella* sp. Pearson (1995) and Pearson and Chaisson (1997) suggested that the ‘chambers’ of *Globigerinatella* that possess areal and/or multiple sutural apertures are homologous with the bullae of *Globigerinita* spp., and that *Globigerinita* was the ancestral form (see also the ontogenetic studies of Bronnimann, 1951, and comments in Bolli and Saunders, 1985). The wall texture of *Globigerinatella* shows a typical microperforate structure identical to that seen in *Globigerinita* and *Tenuitella* (Pearson, 1995; Pearson and Wade, 2009). The evolution of *Globigerinatella* involved the development of highly swollen bullae which, critically, themselves have bullae superimposed upon them, the process being potentially repeated several times during the ontogeny of a single individual. The early growth stage is essentially identical to *Globigerinita*, such that it is only in the adult form that the diagnostic characters become clear. This being the case, designation of an individual to the genus *Globigerinatella* requires the presence of at least one additional bulla superimposed on the bulla-like chamber that is typical of adult *Globigerinita* (see also Bolli and Saunders, 1985, p. 189).

In evolutionary time, there appears to have been a trend in some *Globigerinita* toward more globular test shapes with large inflated bullae that eventually resulted in forms with more than one superimposed bulla; these are the first *Globigerinatella* sp. Only later did areal apertures evolve (the first *G. insueta sensu stricto*), initially as single apertures on the chamber wall (Pearson, 1995), then increasing in number and organization until over 60 apertures can be counted on the final chamber. The same trend has been observed in

both the tropical Pacific Ocean (Chaisson and Leckie, 1993; Pearson, 1995) and tropical Atlantic Ocean (Pearson and Chaisson, 1997). The ‘calibration’ for the first occurrence of *Globigerinatella insueta* in BKSA95 was given as 18.8 Ma, although this was “inferred inasmuch as there is no direct calibration for the FAD of *G. insueta* at present”, nor were BKSA95 aware of the gradual evolution described above.

Of the sites that have so far shown the full evolutionary lineage, the Atlantic Ocean site (Ceara Rise) provides the best opportunity for calibrating the successive first appearances of *Globigerinatella* sp. and *G. insueta sensu stricto*, although it is unfortunate that no magnetostratigraphy is available for them. Pearson and Chaisson (1997) calibrated the events at 20.2 Ma and 17.4 Ma respectively, based on interpolation between other foraminifera and nannofossil datums in the age models for ODP Sites 925 and 929. Of the two *Globigerinatella* FADs, the most useful for re-defining the M2/M3 Zone boundary (=N5/N6 Zone boundary in older schemes) is *Globigerinatella* sp., as there is only a short interval of time (~50 kyr) between the FAD of *G. insueta sensu stricto* and the LAD of *Catapsydrax dissimilis* which marks the M3/M4 (=N6/N7) Zone boundary (see also Pearson, 1995). We therefore revise the zonal definition of Zone M3 to utilize the LO of *Globigerinatella* sp.

Zone M3. *Globigerinatella* sp. / *Catapsydrax dissimilis* Concurrent-range Zone

(herein defined, equivalent to Zone M3 [*Globigerinatella insueta*/ *Catapsydrax dissimilis* Concurrent-range Zone] and upper part of Zone M2 [*Catapsydrax dissimilis* Partial-range Zone] of BKSA95).

941 *Definition:* Concurrent range of the nominate taxon between the LO of *Globigerinatella*
 942 sp. and the HO of *Catapsydrax dissimilis*.

943 *Magnetostratigraphic calibration:* Chron C6n-Chron C5Dr.

944 *Astronomical cycle calibration:* 48_{Mi}-C6n - 44_{Mi}-C5Dn.

945 *Estimated age:* 19.66–17.62 Ma (as per Cande and Kent, 1995); 19.30-17.54 Ma (as per
 946 Lourens et al., 2004); early Miocene.

947 *Remarks:* The new concept of Zone M3 follows from developments in the understanding
 948 of the *Globigerinatella* lineage (discussed above). Within the *Globigerinatella* sp./
 949 *Catapsydrax dissimilis* Concurrent-range Zone (Zone M3) the LO of *Globigerinatella*
 950 *insueta* has been revised from ~18.8 Ma (BKSA95) to 17.69 Ma (interpolated from
 951 Pearson and Chaisson, 1997). The younger age of the LO of *G. insueta* is supported by
 952 studies from Site 1148 (South China Sea, Li et al., 2004), though unfortunately there is
 953 no magnetostratigraphy through this interval.

954

955 **Zone M2. *Globoquadrina binaiensis* Partial-range Zone** (herein amended = lower part
 956 of Zone M2 [*Catapsydrax dissimilis* Partial-range Zone] of BKSA95).

957 *Definition:* Partial-range of the nominate taxon between the HO of ‘*Paragloborotalia*’
 958 *kugleri* and the LO of *Globigerinatella* sp.

959 *Magnetostratigraphic calibration:* Chron C6AAn-Chron C6n.

960 *Astronomical cycle calibration:* 53_{Mi}-C6Ar - 48_{Mi}-C6n.

961 *Estimated age:* 21.81–19.66 Ma (as per Cande and Kent, 1995); 21.12-19.30 Ma (as per
 962 Lourens et al., 2004); early Miocene.

963 *Remarks:* The new concept of Zone M2 follows from developments in the understanding

of the *Globigerinatella* lineage (discussed above). We provisionally refer to *kugleri* and *pseudokugleri* as ‘*Paragloborotalia*’ pending further investigations of these taxa (see Pearson and Wade, 2009 for discussion).

Zone M1. ‘*Paragloborotalia*’ *kugleri* Taxon-range Zone (herein renamed = Zone M1 [*Globorotalia kugleri* Total Range Zone] of BKSA95).

Definition: Total range of the nominate taxon.

Magnetochronologic calibration: Subchron C6Cn.2n-Chron C6AAn.

Astronomical cycle calibration: 57_{Mi}-C6Cn - 53_{Mi}-C6Ar.

Estimated age: 23.73–21.81 Ma (as per Cande and Kent, 1995); 22.96–21.12 Ma (as per Lourens et al., 2004); early Miocene.

Remarks: Renamed to accord with assignment of the species *kugleri* to the temporary genus ‘*Paragloborotalia*’.

Subzone M1b. *Globoquadrina dehiscens* / ‘*Paragloborotalia*’ *kugleri* Concurrent-range Subzone (herein renamed = Subzone M1b [*Globorotalia kugleri*/*Globoquadrina dehiscens* Concurrent-range Subzone] of BKSA95).

Definition: Biostratigraphic interval between the LO of *Globoquadrina dehiscens* and the HO of ‘*Paragloborotalia*’ *kugleri*.

Magnetochronologic calibration: Chron C6Br-Chron C6AAn.

Astronomical cycle calibration: 56_{Mi}-C6Bn - 53_{Mi}-C6Ar.

Estimated age: 23.20–21.81 Ma (as per Cande and Kent, 1995); 22.44–21.12 Ma (as per Lourens et al., 2004); early Miocene.

Remarks: Renamed to accord with assignment of the species *kugleri* to the temporary genus ‘*Paragloborotalia*’.

Subzone M1a. ‘*Paragloborotalia*’ *kugleri* Lowest-occurrence Subzone (herein renamed = Subzone M1a [*Globigerinoides primordius* Interval Subzone] of BKSA95).

Definition: Biostratigraphic interval between the LO of ‘*Paragloborotalia*’ *kugleri* and the LO of *Globoquadrina dehiscens*.

Magnetostratigraphic calibration: Subchron C6Cn.2n-Chron C6Br.

Astronomical cycle calibration: $^{57}\text{Mi-C6Cn} - ^{56}\text{Mi-C6Bn}$.

Estimated age: 23.73–23.20 Ma (as per Cande and Kent, 1995); 22.96–22.44 Ma (as per Lourens et al., 2004); early Miocene.

Remarks: Renamed to accord with assignment of the species *kugleri* to the genus ‘*Paragloborotalia*’.

3.3. Oligocene/Miocene Boundary

The closest planktonic foraminiferal biostratigraphic event to the Oligocene/Miocene Boundary is the LO of ‘*Paragloborotalia*’ *kugleri*. Our recalibrated age from Ceara Rise (Pearson and Chaisson, 1997) of FAD ‘*P*’. *kugleri* is 23.73 Ma. This is 70 kyr younger than the previously published age in BKSA95 but is very consistent with studies from the Lemme-Carrosio Section, where the LO of ‘*P*’. *kugleri* is within Subchron C6Cn.2n (Steininger et al., 1997). Thus the uppermost Oligocene Zone O7 (this study; Zone O6 as per Berggren and Pearson, 2005) extends into the Miocene and reflects the short

stratigraphic interval between the base of the Miocene as designated by the base of
Subchron C6Cn.2n and the LO of '*P. kugleri*.

3. Paleogene Period

For the Paleogene Period all zonal concepts have remained consistent with Berggren and
Pearson (2005), except Zone O6 which we have amended and added Zone O7. The
adjustments to the Paleogene magnetobiochronology are minor and mainly exhibit
revised magnetostratigraphic calibrations. We have updated the calibrated ages of
Paragloborotalia opima and *Chiloguembelina cubensis* as per Wade et al. (2007) which
have slightly modified the duration of Biochrons O5 and O6. In addition we provide
datum events calibrated to the ATS of Pälike et al. (2006) to Zone E11. We have not
attempted to incorporate tuned ages from Ceara Rise for the Oligocene (Pearson and
Chaisson, 1997; Shackleton et al., 1999), because of the significant differences in the age
estimates between Site 925 and 929 (Shackleton et al., 1999, p. 1926). Substantial
revisions occur in the early-mid Eocene (see below).

3.1. Amendment to the Oligocene O Zones

Zone O7. '*Paragloborotalia*' pseudokugleri Lowest-occurrence Zone (herein defined =
upper part of Zone O6 [*Globigerina ciperoensis* Partial-range Zone] of Berggren
and Pearson, 2005).

Definition: Biostratigraphic interval between the LO of the nominate taxon

'Paragloborotalia' pseudokugleri and the LO of '*Paragloborotalia*' *kugleri*.

Magnetochronologic calibration: Subchron C8n.1n-Subchron C6Cn.2n.

1033 *Astronomical cycle calibration: $^{63}\text{Ol-C7Ar}$ - $^{57}\text{Mi-C6Cn}$.*

1034 *Estimated age: 25.9–23.73 Ma (as per Cande and Kent, 1995); 25.4–22.96 Ma (as per*
 1035 *Gradstein et al., 2004); 25.2 (as per Pälike et al., 2006) -22.96 Ma (as per Lourens et al.,*
 1036 *2004); late Oligocene-earliest Miocene.*

1037 *Remarks: The recalibration of the LAD *Paragloborotalia opima* from 27.1 Ma*
 1038 *(BKSA95) to 27.5 Ma (Wade et al., 2007) has resulted in an increase in the duration of*
 1039 *Biochron O6 as per Berggren and Pearson (2005) from 3.3 to 3.7 myr. The FAD of ‘P’.*
 1040 **pseudokugleri* allows greater resolution for this interval. We have subdivided the interval*
 1041 *between HO *P. opima* and LO ‘P’. *kugleri*, using the LO of ‘P’. *pseudokugleri* resulting*
 1042 *in a shorter duration of Biochron O6 and the introduction of Zone O7.*

1043

1044 *The age of the FAD of ‘*Paragloborotalia*’ *pseudokugleri* in BKSA95 is derived from*
 1045 *Hole 803D and Hole 628A (Leckie et al., 1993), where this bioevent was recorded within*
 1046 *Chron C8n and Chron C7n, respectively. Further support for the stratigraphic utility of*
 1047 *‘P’. *pseudokugleri* come from sites drilled during ODP Leg 115 (Premoli Silva and*
 1048 *Spezzaferri, 1990), ODP Leg 208 (Shipboard Scientific Party, 2004) and Site 1148 (Li et*
 1049 *al., 2004); unfortunately, these sites either do not have magnetostratigraphy through this*
 1050 *interval or the magnetostratigraphy is ambiguous. Although this taxon is rare at Site 1218*
 1051 *(equatorial Pacific Ocean), the LO of ‘P’. *pseudokugleri* at 121.56 ± 0.61 meters*
 1052 *composite depth is within Subchron C8n.1n (Shipboard Scientific Party, 2002),*
 1053 *consistent with the age estimate in BKSA95.*

1054

1055 **Zone O6. *Globigerina ciperoensis* Partial-range Zone** (herein amended = lower part of

1056 Zone O6 [*Globigerina ciperoensis* Partial-range Zone] of Berggren and Pearson,
1057 2005).

1058 *Definition:* Biostratigraphic interval characterized by the partial-range of the nominate
1059 taxon, between the HO of *Paragloborotalia opima* and the LO of '*Paragloborotalia*'
1060 *pseudokugleri*.

1061 *Magnetochronologic calibration:* Chron C9n-Subchron C8n.1n.

1062 *Astronomical cycle calibration:* $^{67}\text{Ol-C9n}$ - $^{63}\text{Ol-C7Ar}$.

1063 *Estimated age:* 27.5–25.9 Ma (as per Cande and Kent, 1995); 27.3-25.4 Ma (as per
1064 Luterbacher et al., 2004); 26.9-25.2 Ma (as per Pälike et al., 2006); late Oligocene.

1065 *Remarks:* See discussion above.

1066

1067 **Zone O5. *Paragloborotalia opima* Highest-occurrence Zone**

1068 *Definition:* Biostratigraphic interval between the highest common occurrence (HCO) of
1069 *Chiloguembelina cubensis* and the HO of the nominate taxon *Paragloborotalia*
1070 *opima*.

1071 *Magnetochronologic calibration:* Subchron C10n.1n-Chron C9n.

1072 *Astronomical cycle calibration:* $^{70}\text{Ol-C10n}$ - $^{67}\text{Ol-C9n}$.

1073 *Estimated Age:* 28.4–27.5 Ma (as per Cande and Kent, 1995); 28.3-27.3 Ma (as per
1074 Luterbacher et al., 2004); 28.0-26.9 Ma (as per Pälike et al., 2006); late Oligocene.

1075 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1076 and Pearson (2005). Coccioni et al. (2008) have confirmed that the HCO of *C. cubensis* is
1077 a robust stratigraphic marker for the Oligocene. The age of the LAD of *P. opima* and
1078 LAD of *C. cubensis* have been revised as per Wade et al. (2007).

1079

1080 **Zone O4. *Globigerina angulisuturalis* / *Chiloguembelina cubensis* Concurrent-range**1081 **Zone**1082 *Definition:* Concurrent range of the nominate taxa between the LO of *Globigerina*1083 *angulisuturalis* and the HCO of *Chiloguembelina cubensis*.1084 *Magnetostratigraphic calibration:* Subchron C11n.1n-C10n.1n.1085 *Astronomical cycle calibration:* 73_{Ol-C10r} - 70_{Ol-C10n}.1086 *Estimated age:* 29.4–28.4 Ma (as per Cande and Kent, 1995); 29.5–28.3 Ma (as per

1087 Luterbacher et al., 2004); 29.2–28.0 Ma (as per Pälike et al., 2006); early Oligocene.

1088 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren1089 and Pearson (2005). The FAD of *Globigerina angulisuturalis* has been recorded in Chron

1090 C11n.1n in DSDP Holes 516F (Rio Grande Rise, South Atlantic Ocean; Berggren et al.,

1091 1985) and Site 558 (North Atlantic Ocean; Miller et al., 1985) and (somewhat

1092 equivocally) in C11n.1n in ODP Holes 628A (Little Bahama Bank, North Atlantic

1093 Ocean) and 803D (Ontong Java Plateau, western Pacific Ocean; Leckie et al., 1993; see

1094 also BKSA95; p. 173, Table 10). More recently in the Contessa, Monte Cagnero and

1095 Pieve d'Accinelli sections in the Umbria-Marche Basin, Italy, Coccioni et al. (2008) have

1096 recorded the FAD of *G. angulisuturalis* approximately 400 kyr younger, within mid-

1097 Chron C10r. The discrepancy may lie in preservational bias or diachrony between the

1098 open ocean and Tethys. We retain the calibration of C11n.1n (BKSA95) in this work

1099 pending further investigations.

1100

1101 **Zone O3. *Dentoglobigerina sellii* Partial-range Zone**

- 1102 *Definition:* Partial range of the nominate taxon between the HO of
 1103 *Turborotalia ampliapertura* and the LO of *Globigerina angulisuturalis*.
 1104 *Magnetostratigraphic calibration:* Chron C11r-Subchron C11n.1n.
 1105 *Astronomical cycle calibration:* $^{76}\text{Ol-C11r}$ - $^{73}\text{Ol-C10r}$.
 1106 *Estimated age:* 30.3–29.4 Ma (as per Cande and Kent, 1995); 30.4–29.5 Ma (as per
 1107 Luterbacher et al., 2004); 30.3–29.2 Ma (as per Pälike et al., 2006); early Oligocene.
 1108 *Remarks:* The nomenclature and definition of the zone remains the same as in BKSA95
 1109 and Berggren and Pearson (2005).
 1110
 1111 **Zone O2. *Turborotalia ampliapertura* Highest-occurrence Zone**
 1112 *Definition:* Biostratigraphic interval between the HO of *Pseudohastigerina*
 1113 *nagewichiensis* and the HO of the nominate taxon *Turborotalia ampliapertura*.
 1114 *Magnetostratigraphic calibration:* Chron C12r- Chron C11r.
 1115 *Astronomical cycle calibration:* $^{80}\text{Ol-C12r}$ - $^{76}\text{Ol-C11r}$.
 1116 *Estimated age:* 32.0–30.3 Ma (as per Cande and Kent, 1995); 32.2–30.4 Ma (as per
 1117 Luterbacher et al., 2004); 32.0–30.3 Ma (as per Pälike et al., 2006); early Oligocene.
 1118 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
 1119 and Pearson (2005).
 1120
 1121 **Zone O1. *Pseudohastigerina nagewichiensis* Highest-occurrence Zone**
 1122 *Definition:* Biostratigraphic interval between HO of *Hantkenina alabamensis* and HO of
 1123 the nominate taxon *Pseudohastigerina nagewichiensis* (Berggren and Pearson, 2005).
 1124 *Magnetostratigraphic calibration:* Chron C13r-Chron C12r.

1125 *Astronomical cycle calibration: $^{84}\text{Ol-C13n}$ - $^{80}\text{Ol-C12r}$.*

1126 *Estimated age: 33.7–32.0 Ma (as per Cande and Kent, 1995); 33.9–32.2 Ma (as per*

1127 *Luterbacher et al., 2004); 33.8–32.0 Ma (as per Pälike et al., 2006); early Oligocene.*

1128 *Remarks: The nomenclature and definition of the zone remains the same as in Berggren*

1129 *and Pearson (2005).*

1130

1131 *3.2. Amendment to the Eocene E Zones*

1132 **Zone E16. *Hantkenina alabamensis* Highest-occurrence Zone**

1133 *Definition: Partial range of the nominate taxon between the HO of *Globigerinatheka**

1134 *index and the HO of *Hantkenina alabamensis*.*

1135 *Magnetostratigraphic calibration: Chron C13r–Chron C12r.*

1136 *Astronomical cycle calibration: $^{86}\text{Eo-C13r}$ - $^{84}\text{Ol-C13n}$.*

1137 *Estimated age: 34.3–33.7 Ma (as per Cande and Kent, 1995); 34.5–33.9 Ma (as per*

1138 *Luterbacher et al., 2004); 34.5–33.8 Ma (as per Pälike et al., 2006); late Eocene.*

1139 *Remarks: The nomenclature and definition of the zone remains the same as in Berggren*

1140 *and Pearson (2005). We have added the HCO of *Pseudohastigerina micra* as a secondary*

1141 *marker for the Eocene/Oligocene boundary. *Pseudohastigerina micra* is common in*

1142 *upper Eocene sediments, and this form undergoes a significant size decrease coeval with*

1143 *the extinction of *Hantkenina* (Wade and Pearson, 2008; Wade and Olsson, 2009). This*

1144 *event appears to be coeval between the Indian Ocean (Wade and Pearson, 2008) and the*

1145 *Gulf of Mexico (Miller et al., 2008).*

1146

1147 **Zone E15. *Globigerinatheka index* Highest-occurrence Zone**

1148 *Definition:* Biostratigraphic interval between the HO of *Globigerinatheka semiinvoluta*
 1149 and the HO of the nominate taxon *Globigerinatheka index*.

1150 *Magnetostratigraphic calibration:* Subchron C16n.2n-Chron C13r.

1151 *Astronomical cycle calibration:* $^{89}\text{Eo-C16n}$ - $^{86}\text{Eo-C13r}$.

1152 *Estimated age:* 35.8-34.3 Ma (as per Cande and Kent, 1995); 35.8-34.5 Ma (as per
 1153 Luterbacher et al., 2004); 35.8-34.5 Ma (as per Pälike et al., 2006); late Eocene.

1154 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
 1155 and Pearson (2005).

1156

1157 **Zone E14. *Globigerinatheka semiinvoluta* Highest-occurrence Zone**

1158 *Definition:* Biostratigraphic interval between the HO of *Morozovelloides crassatus* and
 1159 the HO of the nominate taxon, *Globigerinatheka semiinvoluta*.

1160 *Magnetostratigraphic calibration:* Subchron C17n.3n- Subchron C16n.2n.

1161 *Astronomical cycle calibration:* $^{95}\text{Eo-C17n}$ - $^{89}\text{Eo-C16n}$.

1162 *Estimated age:* 38.0-35.8 Ma (as per Cande and Kent, 1995); 37.7-35.8 Ma (as per
 1163 Luterbacher et al., 2004); 38.1-35.8 Ma (as per Pälike et al., 2006); middle-late Eocene.

1164 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
 1165 and Pearson (2005).

1166

1167 **Zone E13. *Morozovelloides crassatus* Highest-occurrence Zone**

1168 *Definition:* Biostratigraphic interval between the HO of *Orbulinoides*
 1169 *beckmanni* and the HO of the nominate taxon, *Morozovelloides crassatus*.

1170 *Magnetostratigraphic calibration:* Subchron C18n.2n-Subchron C17n.3n.

- 1171 *Astronomical cycle calibration: 100_{Eo-C18r} - 95_{Eo-C17n}.*
- 1172 *Estimated age: 40.0-38.0 Ma (as per Cande and Kent, 1995); 39.4-37.7 Ma (as per*
 1173 *Luterbacher et al., 2004); 40.0-38.1 Ma (as per Pälike et al., 2006); middle Eocene.*
- 1174 *Remarks: The nomenclature and definition of the zone remains the same as in Berggren*
 1175 *and Pearson (2006).*
- 1176
- 1177 **Zone E12. *Orbulinoides beckmanni* Taxon-range Zone**
- 1178 *Definition: Total range of the nominate taxon between its LO and HO.*
- 1179 *Magnetostratigraphic calibration: Chron C18r- Subchron C18n.2n.*
- 1180 *Astronomical cycle calibration: 102_{Eo-C18r} - 100_{Eo-C18r}.*
- 1181 *Estimated age: 40.5-40.0 Ma (as per Cande and Kent, 1995); 39.8-39.4 Ma (as per*
 1182 *Luterbacher et al., 2004); 40.8-40.0 Ma (as per Pälike et al., 2006); middle Eocene.*
- 1183 *Remarks: The nomenclature and definition of the zone remains the same as in Berggren*
 1184 *and Pearson (2005). Note there are significant differences in the duration of this biochron*
 1185 *depending upon which time scale is used. Recent studies have suggested that the LO of*
 1186 **Orbulinoides beckmanni* is diachronous (K. Edgar pers.comm.) and the HO in low*
 1187 *latitudes is younger than previous calibrations (Luciani et al., 2010).*
- 1188
- 1189 **Zone E11. *Morozovelloides lehneri* Partial-range Zone**
- 1190 *Definition: Biostratigraphic interval characterized by the partial range of the nominate*
 1191 *taxon between the HO of *Guembelitrioides nuttalli* and the LO of *Orbulinoides**
 1192 **beckmanni*.*
- 1193 *Magnetostratigraphic calibration: Chron C19r-Chron C18r.*

1194 *Estimated age*: 42.3-40.5 Ma (as per Cande and Kent, 1995); 41.4-39.8 Ma (as per
 1195 Luterbacher et al., 2004); middle Eocene.
 1196 *Remarks*: The nomenclature and definition of the zone remains the same as in Berggren
 1197 and Pearson (2006). The age of the HO of *Guembelitrionides nuttalli* is not well
 1198 constrained and requires further study.

1199

1200 **Zone E10. *Acarinina topilensis* Partial-range Zone**

1201 *Definition*: Partial range of the nominate taxon between the HO of
 1202 *Morozovella aragonensis* and the HO of *Guembelitrionides nuttalli*
 1203 *Magnetostratigraphic calibration*: Chron C20n-Chron C19r.
 1204 *Estimated age*: 43.6-42.3 Ma (as per Cande and Kent, 1995); 42.6-41.4 Ma (as per
 1205 Luterbacher et al., 2004); middle Eocene.
 1206 *Remarks*: The nomenclature and definition of the zone remains the same as in Berggren
 1207 and Pearson (2005).

1208

1209 **3.2.1. Early-middle Eocene**

1210 Owing to the pervasive occurrence of chert in the early and middle Eocene (Muttoni and
 1211 Kent, 2007), the interval corresponding to 50-44 Ma (late Ypresian-early Lutetian) and
 1212 the lower/middle Eocene boundary has often proved difficult to recover in deep sea cores.
 1213 This has significantly hampered the correlations of planktonic foraminiferal bioevents to
 1214 the GPTS through this interval. The basal zone of the middle Eocene has traditionally
 1215 been recognized by the lowest occurrence of *Hantkenina* (Bolli, 1957a, b, 1966;
 1216 Berggren et al., 1985, BKS95) which was calibrated by Lowrie et al. (1982) to within

Chron C22n. Due to inconsistencies between the nannofossil and planktonic foraminiferal biostratigraphies, Pearson et al. (2004) suggested that the initial appearance of *Hantkenina* was diachronous. Therefore Berggren and Pearson (2005) used the LO of *Guembelitrionides nuttalli* as the base of Zone E8 corresponding to the lower/middle Eocene (Ypresian-Lutetian) boundary.

In a detailed magnetobiostratigraphic study of an expanded lower-middle Eocene succession from the western Pyrenees, Payros et al. (2007) examined the divergence of planktonic foraminiferal Zones P9 and P10 and the standard zonation (BKSA95). Unlike BKSA95 which places the boundary between planktonic foraminiferal Zones P9 and P10 within calcareous nannofossil Zone NP14 (Subzone CP12a) and magnetic polarity Chrons C22n and C21r, they found the boundary to occur within Zone NP15 (= Zone CP13) and Chron C20r. This is consistent with the biostratigraphic results of Pearson et al. (2004) from Tanzania. Rögl and Egger (2010) have recently corroborated this by recognizing the evolutionary transition of *Clavigerinella* to *Hantkenina* in the upper part of Zone NP15b. The P9/P10 zonal boundary is therefore a surprising 3.1 myr younger than in BKSA95. The significantly younger calibrations for the LO of *Hantkenina* and *Guembelitrionides nuttalli* are also confirmed by Ortiz et al. (2008) and Larrasoana et al. (2008) from the Agost section in Spain. Payros et al. (2007) used the Luterbacher et al. (2004) time scale, here we recalibrate the FAD of *Globigerinatheka kugleri*, *Hantkenina singanoae*, *Guembelitrionides nuttalli* and *Turborotalia frontosa* to Cande and Kent (1995), resulting in FADs of 44.4, 44.5, 46.4 and 49 Ma, respectively (Fig. 2c, Fig. 6, Table 1). This has major implications for the durations of Biochrons E7, E8 and E9 (Fig.

2c).

The revised FAD of *H. singanoae* (Payros et al., 2007) is younger than the previously accepted age of the FAD of *G. kugleri* (as per BKSA95). However, in Tanzania Drilling Project Site 2 and 20 (Nicholas et al., 2006) a short interval exists between these events, equivalent to about 100 kyr (Fig. 6) and suggests a revised age of ~44.4 for the FAD of *G. kugleri*.

The changes in the biochronology through this interval has resulted in the duration of Biochron E7 changing from 1.4 to 4 myr. Payros et al., (2007, 2009) point out the stratigraphic utility of the LO of *Turborotalia frontosa*. Following the earlier zonation of Blow (1979), we have used this event to subdivide Zone E7 into Subzones E7a and E7b (Fig. 2c, Fig. 4c, Fig. 6). Note the E7a/E7b subzonal boundary (this study) is not the same criterion as the E7/E8 zonal boundary of Berggren and Pearson (2005), although on Cande and Kent (1995) magnetostratigraphy, our age for the E7a/E7b subzonal boundary is similar to the value of the E7/E8 zonal boundary in Berggren and Pearson (2005).

There is at present a proposal to define/place the GSSP for the base of the Lutetian Stage (base middle Eocene) at a level equivalent the LO of *Blackites inflatus* (mid-Zone NP14=CP12a/b boundary) within Chron C21r ~47.76 Ma (as per Gradstein et al., 2004) at the Gorronatxe section, Biscaye Province, Spain (Molina et al., in press). This level is younger than the FAD of *Turborotalia frontosa* and within sub-biochron E7b.

1263 **Zone E9. *Globigerinatheka kugleri* / *Morozovella aragonensis* Concurrent-range**

1264 **Zone**

1265 *Definition:* Concurrent range of the nominate taxa between the LO of *Globigerinatheka*
1266 *kugleri* and the HO of *Morozovella aragonensis*.

1267 *Magnetostratigraphic calibration:* Chron C20r (estimated)-Chron C20n.

1268 *Estimated age:* 44.4-43.6 Ma (as per Cande and Kent, 1995); 43.4-42.6 Ma (as per
1269 Luterbacher et al., 2004); middle Eocene.

1270 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1271 and Pearson (2005). The age of the LO of *Globigerinatheka kugleri* is not well
1272 constrained and there is a significant need for further study of this interval in continuous
1273 sections with magnetostratigraphy (see discussion above).

1274

1275 **Zone E8. *Guembelitrionides nuttalli* Lowest-occurrence Zone**

1276 *Definition:* Biostratigraphic interval between the LO of the nominate taxon
1277 *Guembelitrionides nuttalli* and the LO of *Globigerinatheka kugleri*.

1278 *Magnetostratigraphic calibration:* Chron C21n-Chron C20r (estimated).

1279 *Estimated age:* 46.4-44.4 Ma (as per Cande and Kent, 1995); 45.5-43.4 Ma (as per
1280 Luterbacher et al., 2004); middle Eocene.

1281 *Remarks:* While the nomenclature and definition of the zone remains the same as in
1282 Berggren and Pearson (2005), the duration and boundary ages of the biochron have
1283 changed significantly (see discussion above). Payros et al. (2007) find the LO of
1284 *Guembelitrionides nuttalli* in upper Chron C21n. However, a slightly younger level within
1285 Chron C20r was found by Larrasoana et al. (2008).

1286

1287 **Zone E7. *Acarinina cuneicamerata* Lowest-occurrence Zone**

1288 *Definition:* Biostratigraphic interval between the LO of the nominate taxon *Acarinina*
 1289 *cuneicamerata* and the LO of *Guembelitrioides nuttalli*.

1290 *Magnetostratigraphic calibration:* Chron C22r (estimated) to Chron C21n.

1291 *Estimated age:* 50.4–46.4 Ma (as per Cande and Kent, 1995); 50.3–45.5 Ma (as per
 1292 Luterbacher et al., 2004); early-middle Eocene.

1293 *Remarks:* The definition of this zone remains the same as in Berggren and Pearson
 1294 (2005), however, the estimated duration of the biochron is much longer because of
 1295 recalibration of the LO of *Guembelitrioides nuttalli* (Payros et al., 2007).

1296

1297 **Subzone E7b. *Turborotalia frontosa* Lowest-occurrence Subzone**

1298 *Definition:* Biostratigraphic interval between the LO of the nominate taxon *Turborotalia*
 1299 *frontosa* and the LO of *Guembelitrioides nuttalli*.

1300 *Magnetostratigraphic calibration:* Chron C21r to Chron C21n.

1301 *Estimated age:* 49.0–46.4 Ma (as per Cande and Kent, 1995); 48.6–45.5 Ma (as per
 1302 Luterbacher et al., 2004); middle Eocene.

1303 *Remarks:* Zone E7 has been subdivided based on the LO of *Turborotalia frontosa*
 1304 (Payros et al., 2007). See discussion in Berggren and Miller (1988) regarding previous
 1305 use of *T. frontosa* as a zonal marker

1306

1307 **Subzone E7a. *Acarinina cuneicamerata* Lowest-occurrence Subzone**

1308 *Definition:* Biostratigraphic interval between the LO of the nominate taxon *Acarinina*

- 1309 *cuneicamerata* and the LO *Turborotalia frontosa*.
- 1310 *Magnetostratigraphic calibration*: Chron C22r-Chron C21r.
- 1311 *Estimated age*: 50.4–49.0 Ma (as per Cande and Kent, 1995); 50.3–48.6 Ma (as per
- 1312 Luterbacher et al., 2004); early Eocene.
- 1313 *Remarks*: Zone E7 has been subdivided based on the LO of *Turborotalia frontosa*
- 1314 (Payros et al., 2007).
- 1315
- 1316 **Zone E6. *Acarinina pentacamerata* Partial-range Zone**
- 1317 *Definition*: Partial range of the nominate taxon between the HO of *Morozovella*
- 1318 *subbotinae* and the LO of *Acarinina cuneicamerata*.
- 1319 *Magnetostratigraphic calibration*: Subchron 23n.1n-Chron C22r.
- 1320 *Estimated age*: 50.8–50.4 Ma (as per Cande and Kent, 1995); 50.8–50.3 Ma (as per
- 1321 Luterbacher et al., 2004); early Eocene.
- 1322 *Remarks*: The nomenclature and definition of the zone remains the same as in Berggren
- 1323 and Pearson (2005).
- 1324
- 1325 **Zone E5. *Morozovella aragonensis* / *Morozovella subbotinae* Concurrent-range Zone**
- 1326 *Definition*: Concurrent range of the nominate taxa between the LO of *Morozovella*
- 1327 *aragonensis* and the HO of *Morozovella subbotinae*.
- 1328 *Magnetostratigraphic calibration*: Chron C23r-Subchron C23n.1n.
- 1329 *Estimated age*: 52.3–50.8 Ma (as per Cande and Kent, 1995 and Luterbacher et al., 2004);
- 1330 early Eocene.
- 1331 *Remarks*: The nomenclature and definition of the zone remains the same as in Berggren

1332 and Pearson (2005).

1333

1334 **Zone E4. *Morozovella formosa* Lowest-occurrence Zone**

1335 *Definition:* Biostratigraphic interval between the LO of the nominate taxon *Morozovella*
1336 *formosa* and the LO of *Morozovella aragonensis*.

1337 *Magnetostratigraphic calibration:* Chron C24r-Chron C23r.

1338 *Estimated age:* 54.0-52.3 Ma (as per Cande and Kent, 1995); 54.4-52.3 Ma (as per
1339 Luterbacher et al., 2004); early Eocene.

1340 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1341 and Pearson (2005).

1342

1343 **Zone E3. *Morozovella marginodentata* Partial-range Zone**

1344 *Definition:* Biostratigraphic interval characterized by the partial range of the nominate
1345 taxon between the HO of *Morozovella velascoensis* and LO of *Morozovella formosa*.

1346 *Magnetostratigraphic calibration:* Chron C24r.

1347 *Estimated age:* 54.5-54.0 Ma (as per Cande and Kent, 1995); 54.9-54.4 Ma (as per
1348 Luterbacher et al., 2004); early Eocene.

1349 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1350 and Pearson (2005).

1351

1352 **Zone E2. *Pseudohastigerina wilcoxensis* / *Morozovella velascoensis***

1353 **Concurrent-range Zone**

1354 *Definition:* Biostratigraphic interval characterized by the concurrent biostratigraphic

1355 ranges of the nominate taxa between the LO of *Pseudohastigerina wilcoxensis* and the
 1356 HO of *Morozovella velascoensis*.

1357 *Magnetostratigraphic calibration*: Chron C24r.

1358 *Estimated age*: 55.4-54.5 Ma (as per Cande and Kent, 1995); 55.7-54.9 Ma (as per
 1359 Luterbacher et al., 2004); early Eocene.

1360 *Remarks*: The nomenclature and definition of the zone remains the same as in Berggren
 1361 and Pearson (2005).

1362

1363 **Zone E1. *Acarinina sibaiaensis* Lowest-occurrence Zone**

1364 *Definition*: Biostratigraphic interval between the LO of the nominate taxon *Acarinina*
 1365 *sibaiaensis* and the LO of *Pseudohastigerina wilcoxensis*.

1366 *Magnetostratigraphic calibration*: Chron C24r.

1367 *Estimated age*: 55.5-55.4 Ma (as per Cande and Kent, 1995); 55.8-55.7 Ma (as per
 1368 Luterbacher et al., 2004); early Eocene.

1369 *Remarks*: The nomenclature and definition of the zone remains the same as in Berggren
 1370 and Pearson (2005).

1371

1372 *3.3. Amendment to the Paleocene P Zones*

1373 **Zone P5. *Morozovella velascoensis* Partial-range Zone**

1374 *Definition*: Biostratigraphic interval characterized by the partial range of the nominate
 1375 taxon between the HO of *Globanomalina pseudomenardii* and the LO of *Acarinina*
 1376 *sibaiaensis*.

1377 *Magnetostratigraphic calibration*: Chron C25n-Chron C24r.

1378 *Estimated Age:* 55.9-55.5 Ma (as per Cande and Kent, 1995); 56.7-55.8 Ma (as per
 1379 Luterbacher et al., 2004); late Paleocene.
 1380 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
 1381 and Pearson (2005).

1382

1383 **Zone P4. *Globanomalina pseudomenardii* Taxon-range Zone.**

1384 *Definition:* Biostratigraphic interval characterized by the total range of the nominate
 1385 taxon *Globanomalina pseudomenardii*.

1386 *Magnetochronologic calibration:* Chron C26r-Chron C25n.

1387 *Estimated age:* 59.4-55.9 Ma (as per Cande and Kent, 1995); 60.2-56.7 Ma (as per
 1388 Luterbacher et al., 2004); middle-late Paleocene.

1389 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
 1390 and Pearson (2005).

1391

1392 **Subzone P4c. *Acarinina soldadoensis* / *Globanomalina pseudomenardii***

1393 **Concurrent-range Subzone.**

1394 *Definition:* Concurrent range of the nominate taxa from the LO of *Acarinina soldadoensis*
 1395 to the HO of *Globanomalina pseudomenardii*.

1396 *Magnetic calibration:* Chron C25r-Chron C25n.

1397 *Age estimate:* 56.5-55.9 Ma (as per Cande and Kent, 1995); 57.3-56.7 Ma (as per
 1398 Luterbacher et al., 2004); late Paleocene.

1399 *Remarks:* The definition of this subzone remains the same as that of BKSA95.

1400

1401 **Subzone P4b. *Acarinina subsphaerica* Partial-range Subzone**

1402 *Definition:* Partial range of the nominate taxon *Acarinina subsphaerica* from the HO of

1403 *Parasubbotina variospira* to the LO of *Acarinina soldadoensis*.

1404 *Magnetic calibration:* Chron C26r-Chron 25r.

1405 *Age estimate:* 59.2-56.5 Ma (as per Cande and Kent, 1995); 60.0-57.3 Ma (as per

1406 Luterbacher et al., 2004); middle-late Paleocene.

1407 *Remarks:* The definition of this subzone remains the same as that of Berggren et al.

1408 (2000).

1409

1410 **Subzone P4a. *Globanomalina pseudomenardii* / *Parasubbotina variospira***

1411 **Concurrent-range Subzone**

1412 *Definition:* Concurrent range of the nominate taxa from the LO of *Globanomalina*

1413 *pseudomenardii* to the HO of *Parasubbotina variospira*.

1414 *Magnetic calibration:* Chron C26r.

1415 *Age estimate:* 59.4-59.2 Ma (as per Cande and Kent, 1995); 60.2-60.0 Ma (as per

1416 Luterbacher et al., 2004); middle Paleocene.

1417 *Remarks:* The definition of this subzone remains the same as that of Berggren et al.

1418 (2000).

1419

1420 **Zone P3. *Morozovella angulata* Lowest-occurrence Zone**

1421 *Definition:* Biostratigraphic interval between the LO of *Morozovella angulata* and the LO

1422 of *Globanomalina pseudomenardii*.

1423 *Magnetochronologic calibration:* Chron C27n-Chron C26r.

1424 *Estimated age:* 61.0-59.4 Ma (as per Cande and Kent, 1995); 61.7-60.2 Ma (as per
1425 Luterbacher et al., 2004); early-middle Paleocene.
1426 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1427 and Pearson (2005).

1428

1429 **Subzone P3b. *Igorina albeari* Lowest-occurrence Subzone**

1430 *Definition:* Biostratigraphic interval from the LO of *Igorina albeari* to the LO of
1431 *Globanomalina pseudomenardii*.

1432 *Magnetic calibration:* Chron C26r-Chron C26.

1433 *Age estimate:* 60.0-59.4 Ma (as per Cande and Kent, 1995); 60.8-60.2 Ma (as per
1434 Luterbacher et al., 2004); middle Paleocene.

1435 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1436 and Pearson (2005).

1437

1438 **Subzone P3a. *Igorina pusilla* Partial-range Subzone**

1439 *Definition:* Biostratigraphic interval defined by the partial range of *Igorina pusilla*
1440 between the LO of *Morozovella angulata* and the LO of *Igorina albeari*.

1441 *Magnetic calibration:* Chron C27n-Chron 26r.

1442 *Age estimate:* 61.0-60.0 Ma (as per Cande and Kent, 1995); 61.7-60.8 Ma (as per
1443 Luterbacher et al., 2004); early Paleocene.

1444 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1445 and Pearson (2005).

1446

1447 **Zone P2. *Praemurica uncinata* Lowest-occurrence Zone**

1448 *Definition:* Biostratigraphic interval between the LO of *Praemurica uncinata* and the LO
1449 of *Morozovella angulata*.

1450 *Magnetochronologic calibration:* Chron C27r-Chron C27n.

1451 *Estimated age:* 61.4-61.0 Ma (as per Cande and Kent, 1995); 62.1-61.7 Ma (as per
1452 Luterbacher et al., 2004); early Paleocene.

1453 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1454 and Pearson (2005).

1455

1456 **Zone P1. *Eoglobigerina edita* Partial-range Zone**

1457 *Definition:* Partial range of the nominate taxon between the HO of

1458 *Parvularugoglobigerina eugubina* and the LO of *Praemurica uncinata*.

1459 *Magnetochronologic calibration:* Chron C29r-Chron C27r.

1460 *Estimated age:* 64.8-61.4 (as per Cande and Kent, 1995); 65.2-62.1 Ma (as per
1461 Luterbacher et al., 2004); early Paleocene.

1462 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren
1463 and Pearson (2005).

1464

1465 **Subzone P1c. *Globanomalina compressa* Lowest-occurrence Subzone** (herein

1466 amended and renamed = Subzone P1c [*Globanomalina compressa*/*Praemurica*
1467 *inconstans* Lowest-occurrence Subzone] of Berggren and Pearson, 2005; =

1468 Subzone P1c [*Globanomalina compressa*/*Praemurica inconstans*-*Praemurica*
1469 *uncinata* Subzone] of BKSA95).

1470 *Definition:* Biostratigraphic interval between the LO of *Globanomalina compressa* and
1471 the LO of *Praemurica uncinata*.

1472 *Magnetic calibration:* Chron C28n-Chron C27r.

1473 *Age estimate:* 62.9-61.4 (as per Cande and Kent, 1995); 63.5-62.1 Ma (as per Luterbacher
1474 et al., 2004); early Paleocene.

1475 *Remarks:* The definition used here removes the operational ambiguity inherent in the
1476 “and/or” designation in Berggren and Pearson, (2005), in which the Subzone was defined
1477 as the biostratigraphic interval between the LO of *Globanomalina compressa* and/or
1478 *Praemurica inconstans* and the LO of *Praemurica uncinata*.

1479

1480 **Subzone P1b. *Subbotina triloculinoides* Lowest-occurrence Subzone** (herein amended
1481 = Subzone P1b [*Subbotina triloculinoides* Lowest-occurrence Subzone] of
1482 Berggren and Pearson, 2005; = Subzone P1b [*Subbotina triloculinoides*-
1483 *Globanomalina compressa*/*Praemurica inconstans* Interval Subzone] of
1484 BKSA95).

1485 *Definition:* Biostratigraphic interval between the LO of *Subbotina triloculinoides* and the
1486 LO of *Globanomalina compressa*.

1487 *Magnetic calibration:* Chron C29n-Chron C28n.

1488 *Estimated age:* 64.3-62.9 (as per Cande and Kent, 1995); 64.7-63.5 Ma (as per
1489 Luterbacher et al., 2004); early Paleocene.

1490 *Remarks:* The definition used here removes the operational ambiguity inherent in the
1491 “and/or” designation in Berggren and Pearson, (2005), in which the subzone was defined
1492 as the biostratigraphic interval between the LO of *Subbotina triloculinoides* and the LOs

1493 of *Globanomalina compressa* and/or *Praemurica inconstans*.

1494

1495 **Subzone P1a. *Parasubbotina pseudobulloides* Partial-range Subzone**

1496 *Definition:* Partial range of the nominate taxon between the HO of

1497 *Parvularugoglobigerina eugubina* and the LO of *Subbotina triloculinoides*.

1498 *Magnetic calibration:* Chron C29r-Chron C29n.

1499 *Estimated age:* 64.8-64.3 (as per Cande and Kent, 1995); 65.2-64.7 Ma (as per

1500 Luterbacher et al., 2004); early Paleocene.

1501 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren

1502 and Pearson (2005).

1503

1504 **Zone Pα. *Parvularugoglobigerina eugubina* Taxon-range Zone**

1505 *Definition:* Biostratigraphic interval characterized by the total range of the nominate

1506 taxon *Parvularugoglobigerina eugubina*.

1507 *Magnetostratigraphic calibration:* Chron C29r.

1508 *Estimated age:* 64.97-64.8 (as per Cande and Kent, 1995); 65.46-65.2 Ma (as per

1509 Luterbacher et al., 2004); early Paleocene.

1510 *Remarks:* The nomenclature and definition of the zone remains the same as in Berggren

1511 and Pearson (2005).

1512

1513 **Zone P0. *Guembelitria cretacea* Partial-range Zone**

1514 *Definition:* Biostratigraphic interval characterized by the partial range of the nominate

1515 taxon between the HO of Cretaceous taxa (e.g., *Globotruncana*, *Rugoglobigerina*,

Globigerinelloides) and the LO of *Parvularugoglobigerina eugubina*.

Magnetochronologic calibration: Chron C29r.

Estimated age: 65.0-64.97 (as per Cande and Kent, 1995); 65.5-65.46 Ma (as per

Luterbacher et al., 2004); early Paleocene.

Remarks: The nomenclature and definition of the zone remains the same as in Berggren

and Pearson (2005).

4. Conclusions

We compile 187 revised calibrations of planktonic foraminiferal bioevents for the Cenozoic and provide calibrations to the GPTS of the Cenozoic and ATS of the Neogene and late Paleogene. Our compilation provides a template for Cenozoic magnetobio- and magnetobioastro-chronology. With recent progress in astronomical tuning, it is clear that high resolution biostratigraphic work and integrated biochronologies are needed to reduce the uncertainty of a number of events and study potential diachrony between the Atlantic and Pacific oceans. Future developments in radioisotopic and astronomical dating will undoubtedly lead to further revision and refinements in Cenozoic planktonic foraminiferal biochronology.

Acknowledgements

Two anonymous reviewers helped to improve the manuscript. Mark Leckie, Frédéric Quillévéré, Henk Brinkhuis and Eustoquio Molina provided valuable comments and discussion. Our thanks are extended to Tracy Aze for providing the template used in Fig. 2 and the Shipboard Scientific Party of IODP Expedition 320/321 for discussion and the

1539 inspiration for this compilation. BSW acknowledges support from National Science
1540 Foundation (NSF) CAREER Award (EAR-0847300), Consortium for Ocean
1541 Leadership/NSF (OCE- 0352500) and the Natural Environment Research Council
1542 (NE/G014817/1).
1543

References

- Aguirre, E., Pasini, G., 1985. The Pliocene-Pleistocene Boundary. *Episodes* 8, 116-120.
- Backman, J., Raffi, I., 1997. Calibration of Miocene nannofossil events to orbitally-tuned cyclostratigraphies from Ceara Rise. In: Curry, W.B., Shackleton, N.J., Richter, C., Bralower, T.J. (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results*, College Station, TX (Ocean Drilling Program) 154, 83–99.
- Bandy, O.L., 1964. Cenozoic planktonic foraminiferal zonation. *Micropaleontology* 10, 1-17.
- Bandy, O. L., 1972. Origin and development of *Globorotalia (Turborotalia) pachyderma* (Ehrenberg). *Micropaleontology* 18, 294-318.
- Banner, F. T., Blow, W.H., 1965. Progress in the planktonic foraminiferal biostratigraphy of the Neogene. *Nature* 208, 1164-1166.
- Berggren, W.A., 1969. Cenozoic chronostratigraphy, planktonic foraminiferal zonation and the radiometric time scale. *Nature* 224, 1072-1075.
- Berggren, W.A., 1973. The Pliocene Time Scale: calibration of planktonic foraminiferal and calcareous nannoplankton zones. *Nature* 243, 391-397.
- Berggren, W.A. 1977. Late Neogene planktonic foraminiferal biostratigraphy of DSDP Site 357 (Rio Grande Rise). In: Perch-Nielsen et al., *Initial Reports of the Deep Sea Drilling Project XXXIX*, 591 – 614.
- Berggren, W.A., 1993. Neogene planktonic foraminiferal biostratigraphy of eastern Jamaica. In: Wright, R.M., Robinson, (Eds.), *Biostratigraphy of Jamaica*. Geological Society of America Memoir 182, 179-217.

- 1566 Berggren, W.A., Miller, K.G. 1988. Paleogene tropical planktonic foraminiferal
1567 biostratigraphy and magnetobiochronology. *Micropaleontology* 34, 362 – 380.
- 1568 Berggren, W. A., Pearson, P. N. 2005. A revised tropical and subtropical Paleogene
1569 planktonic foraminiferal zonation. *Journal of Foraminiferal Research* 35, 279-298.
- 1570 Berggren, W. A., Pearson, P. N. 2006. Tropical and subtropical planktonic foraminiferal
1571 zonation of the Eocene and Oligocene. In: Pearson, P.N., Olsson, R.K., Huber, B.T.,
1572 Hemleben, C. and Berggren, W.A. (Eds.), *Atlas of Eocene Planktonic Foraminifera*,
1573 Cushman Foundation Special Publication, 41: 29-40.
- 1574 Berggren, W.A., Hamilton, N., Johnson, D.A., Pujol, C., Weiss, W., Weiss, P., Cepek, P.,
1575 Gombos, Jr., A.M., 1983. Magnetobiostratigraphy of Deep Sea Drilling Project Leg
1576 72, Sites 515-518, Rio Grande Rise (South Atlantic). Initial Reports of the Deep Sea
1577 Drilling Project 72, 939 – 948.
- 1578 Berggren, W.A., Kent, D.V., Flynn, J.J., Van Couvering, J.A. 1985. Cenozoic
1579 geochronology. *Geological Society of America Bulletin* 96, 1407–1418.
- 1580 Berggren, W.A., Hilgen, F.J., Langereis, C.G., Kent, D.V., Obradovich, J.D., Raffi, I.,
1581 Raymo, M.E., Shackleton, N.J., 1995a. Late Neogene chronology: new perspectives
1582 in high-resolution stratigraphy. *Geological Society of America Bulletin* 107, 1272–
1583 1287.
- 1584 Berggren, W. A., Kent, D. V., Swisher, C. C. III, Aubry, M.-P., 1995b. A revised
1585 Cenozoic geochronology and chronostratigraphy. In: Berggren, W. A., Kent, D. V.,
1586 Aubry, M.-P., Hardenbol, J. (Eds.), *Geochronology, Time Scales and Global*
1587 *Stratigraphic Correlation: A Unified Temporal Framework for an Historical*
1588 *Geology*, SEPM Spec. Publ., vol. 54, pp. 129 - 212.

- 1589 Berggren, W.A., Aubry, M.-P., van Fossen, M., Kent, D.V., Norris, R.D., Quillévéré, F.,
1590 2000. Integrated Paleocene calcareous plankton magnetobiochronology and stable
1591 isotope stratigraphy: DSDP Site 384 (NW Atlantic Ocean): Palaeogeography,
1592 Palaeoclimatology, Palaeoecology 159, 1–51.
- 1593 Blow, W.H., 1957. Transatlantic correlation of Miocene sediments. Micropaleontology 3,
1594 77-79
- 1595 Blow, W.H., 1959. Age, correlation, and biostratigraphy of the upper Tocuyo (San
1596 Lorenzo) and Pown formations, eastern Falcon, Venezuela. Bulls. Am. Paleontology
1597 39, 67-251.
- 1598 Blow, W. H., 1969. Late Middle Eocene to Recent planktonic foraminiferal
1599 biostratigraphy. In: Brönnimann, P., Renz, H. H., (Eds.), Proceedings of the First
1600 International Conference on Planktonic Microfossils: Leiden: E.J. Brill, 1, 199-422.
- 1601 Blow, W.H., 1979. The Cainozoic Foraminiferida, v. I & 11, Leiden, E. J. Brill, 1413 p.
- 1602 Blow, W.H., Banner, F., 1962. The Mid-Tertiary (Upper Eocene to Aquitanian)
1603 Globigerinaceae. In: Eames F.E et al., (Eds.), Fundamentals of Mid-Tertiary
1604 Stratigraphical Correlation, Cambridge University Press, 61-163.
- 1605 Blow, W.H., Banner, F., 1966. The morphology, taxonomy and biostratigraphy of
1606 *Globorotalia barisanensis* LeRoy, *Globorotalia fohsi* Cushman and Ellisor, and
1607 related taxa. Micropaleontology 12, 286-303.
- 1608 Bolli, H. M., 1957a. The genera *Globigerina* and *Globorotalia* in the Paleocene-Lower
1609 Eocene Lizard Springs Formation of Trinidad, B.W.I., In: Loeblich, A. R., Jr., and
1610 collaborators (Eds.), Studies in Foraminifera: Bulletin of the United States National
1611 Museum 215, 61–82.

- 1612 Bolli, H.M., 1957b. Planktonic foraminifera from the Eocene Navet Formation and San
1613 Fernando Formations in Trinidad, B.W.I. In: Loeblich, A. R., Jr., and collaborators
1614 (Eds.), Studies in Foraminifera: Bulletin of the United States National Museum 215,
1615 155–172.
- 1616 Bolli, H.M., 1966. Zonation of Cretaceous to Pliocene marine sediments based on
1617 planktonic foraminifera: Boletín Informativo Asociación Venezolana de Geología,
1618 Mineraría y Petróleo 9, 3–32.
- 1619 Bolli, H.M., Bermúdez, P.J., 1965. Zonation based on planktonic foraminifera of Middle
1620 Miocene to Pliocene warm-water sediments. Asoc. Venezolana Geología, Minería y
1621 Petróleo Bol. Inf 8, 119-149.
- 1622 Bolli, H. M., Saunders, J. B., 1985. Oligocene to Holocene low latitude planktic
1623 foraminifera. In: Bolli, H. M., Saunders, J. B., Perch-Nielsen, K. (Eds.), Plankton
1624 Stratigraphy, 1, Cambridge University Press, Cambridge, 155 - 262.
- 1625 Brönnimann, P., 1951. *Globigerinita naparimaensis*, n. gen., n. sp., from the Miocene of
1626 Trinidad, B.W.I. Contr. Cushman Founda. Foram, Res. 2, 16-18.
- 1627 Cande, S. C., Kent, D. V., 1995. Revised calibration of the geomagnetic polarity
1628 timescale for the Late Cretaceous and Cenozoic. Journal of Geophysical Research
1629 100, 6093 – 6095.
- 1630 Chaisson, W. P, Leckie, R. M. 1993. High-resolution Neogene planktonic foraminifer
1631 biostratigraphy of Site 806, Ontong Java Plateau (western Equatorial Pacific). In:
1632 Berger, W. H., Kroenke, L. W., Mayer, L. A., et al., Proceedings of the Ocean
1633 Drilling Program, Scientific Results 130, 137 - 178.

- 1634 Chaisson, W. P, Pearson, P. N. 1997. Planktonic foraminifer biostratigraphy at Site 925:
1635 Middle Miocene – Pleistocene. In: Shackleton, N. J., Curry, W. B., Richter, C.,
1636 Bralower, T. J. (Eds.), Proceeding of the Ocean Drilling Program, Scientific Results
1637 154, 3 - 31.
- 1638 Chaproniere, G. C. H., Styzen, M. J., Sager, W. W., Nishi, H. Quinterno, P. J.,
1639 Abrahamsen, N., 1994. Late Neogene biostratigraphic and magnetostratigraphic
1640 synthesis. Proceeding of the Ocean Drilling Program, Scientific Results 135, 857-
1641 877.
- 1642 Cita, M.B., 1973. Pliocene biostratigraphy and chronostratigraphy. In: Ryan, W.F.F. et
1643 al., Initial Reports of the Deep Sea Drilling Project, Washington, D.C., Government
1644 Printing Office 13, 1343-1379.
- 1645 Coccioni, R., Marsili, A., Montanari, A., Bellanca, A., Neri, R., Bice, D.M., Brinkhuis,
1646 H., Church, N., Macalady, A., McDaniel, A., Deino, A., Lirer, F., Sprovieri, M.,
1647 Maiorano, P., Monechi, S., Nini, C., Nocchi, M., Pross, J., Rochette, P., Sagnotti, L.,
1648 Tateo, F., Touchard, Y., Van Simaeys, S., Williams, G.L., 2008. Integrated
1649 stratigraphy of the Oligocene pelagic sequence in the Umbria-Marche basin
1650 (northeastern Apennines, Italy): A potential Global Stratotype Section and Point
1651 (GSSP) for the Rupelian/Chattian boundary. Geological Society of America Bulletin
1652 120, 487–511.
- 1653 Curry, W.B., Shackleton, N.J., Richter, C., et al., 1995. Proceeding of the Ocean Drilling
1654 Program, Initial Reports, 154, College Station, TX (Ocean Drilling Program).
- 1655 Cushman, J.A., Bermúdez, P.J., 1949. Some Cuban species of *Globorotalia*. Cushman
1656 Lab. Foram. Research Contrib. 25, 26-45

- 1657 Cushman, J.A., Stainforth, R.M.S., 1945. The foraminifera of the Cipero Marl Formation
1658 of Trinidad, British West Indies. Cushman Laboratory for Foraminiferal Research.
1659 Special Publication. 14, 75 pp.
- 1660 Dowsett, H.J., 1988. Diachrony of late Neogene microfossils in the Southwest Pacific
1661 Ocean: Application of the graphic correlation method. *Paleoceanography* 3, 209-
1662 222.
- 1663 Gibbard, P. L., Head, M. J., Walker, M. J. C., the Subcommittee on Quaternary
1664 Stratigraphy, 2009. Formal ratification of the Quaternary System/Period and the
1665 Pleistocene Series/Epoch with a base at 2.58 Ma. *J. Quaternary Sci.*, in press.
- 1666 Gradstein, F.M., Ogg, J.G., Smith, A.G., 2009 (Eds.), *A Geologic Time Scale 2004*:
1667 Cambridge University Press, Cambridge, UK, 500 pp.
- 1668 Hancock, H. J. L., Chaproniere, G. C., Dickens, G.R., Henderson, R.A., 2002. Early
1669 Paleocene planktic foraminiferal and carbon isotope stratigraphy, Hole 762C,
1670 Exmouth Plateau, northwest Australian margin. *Journal of Micropalaeontology* 21,
1671 29–42.
- 1672 Hedberg, H.D., (Ed.) 1976. *International stratigraphic guide: a guide to stratigraphic*
1673 *classification, terminology and procedure*. John Wiley and Sons, New York.
- 1674 Hilgen, F.J., Krijgsman, W., Raffi, I., Turco, E., Zachariasse, W.J., 2000. Integrated
1675 stratigraphy and astronomical calibration of the Serravallian/Tortonian boundary
1676 section at Monte Gibliscemi (Sicily, Italy). *Marine Micropaleontology* 38, 181–211.
- 1677 Hilgen, F., Brinkhuis, H., Zachariasse, W.-J. 2006. Unit stratotypes for global stages: The
1678 Neogene perspective. *Earth-Science Reviews* 74, 113-125.
- 1679 Hodell, D.A. and Kennett, J. P., 1986. Late Miocene-early Pliocene stratigraphy and

- 1680 paleoceanography of the South Atlantic and southwest Pacific Oceans: a synthesis.
 1681 Paleoceanography 1, 285-311.
- 1682 Jenkins, D. G., 1966. Planktonic foraminiferal zones and new taxa from the Danian to
 1683 Lower Miocene of New Zealand, New Zealand Jour. Geology and Geophysics 8,
 1684 1088-1126.
- 1685 Jenkins, D. G., 1967. Planktonic foraminiferal zones and new taxa from the lower
 1686 Miocene to the Pleistocene of New Zealand, New Zealand Jour. Geology and
 1687 Geophysics 10, 1064-1078.
- 1688 Jenkins, D. G., 1971. New Zealand Cenozoic planktonic foraminifera, New Zealand
 1689 Geol. Survey Paleont. Bull. 42, 1-278.
- 1690 Jenkins, D.G., Saunders, J.B., and Cifelli, R., 1981. The relationship of *Globigerinoides*
 1691 *bisphericus* Todd 1954 to *Praeorbulina sicana* (de Stefani) 1952.J. Foraminiferal
 1692 Res. 11, 262-267.
- 1693 Kaneps, A. G., 1975. Cenozoic planktonic foraminifera from Antarctic Deep Sea
 1694 sediments, Leg 28, DSDP. In: Hays, D. E. et al., Initial Reports of the Deep Sea
 1695 Drilling Project, Washington, D.C., US. Government Printing Office 28, 573-583.
- 1696 Keigwin, L.D., Jr., 1982. Neogene planktonic foraminifers from Deep Sea Drilling
 1697 Project Sites 502 and 503. Initial Reports of the Deep Sea Drilling Project,
 1698 Washington, D.C., US. Government Printing Office 68, 269-288.
- 1699 Kennett, J.P., Srinivasan, M.S., 1983. Neogene Planktonic Foraminifera , A Phylogenetic
 1700 Atlas. Hutchinson Ross, Stroudsburg, Pennsylvania, 265 pp.
- 1701 Kennett, J.P., Srinivasan, M.S., 1984. Neogene datum planes: foraminiferal successions
 1702 in Australia with reference sections from the Ninetyeast Ridge and the Ontong-Java

- 1703 Plateau. In: Ikebe, N., Tsuchi, R. (Eds.), Pacific Neogene Datum Planes,
1704 Contributions to Biostratigraphy and Chronology: Tokyo (Univ. of Tokyo Press),
1705 11-33.
- 1706 Krasheninikov, V.A., 1971. Cenozoic foraminifera. In: Fischer, A.G., et al., Initial
1707 Reports of the Deep Sea Drilling Project VI, 1055 – 1068.
- 1708 Larrasoana, J.C., Gonzalvo, C., Molina, E., Monechi, S., Ortiz, S., Tori, F. & Tosquella,
1709 J., 2008. Integrated magnetobiochronology of the Early/Middle Eocene transition at
1710 Agost (Spain): implications for defining the Ypresian/Lutetian boundary stratotype.
1711 *Lethaia* 41, 395–415.
- 1712 Laskar, J., 1999. The limits of Earth orbital calculations for geological time-scale use,
1713 *Philos. Trans. R. Soc. London, Ser. A*, 357(1757), 1735– 1759,
1714 doi:10.1098/rsta.1999.0399.
- 1715 Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. Correia, Levrard B., 2004. A long term
1716 numerical solution for the insolation quantities of the Earth. *Astron. Astrophys.*, 428,
1717 261
- 1718 Li, Q., Radford, S.S., Banner, F.T., 1992. Distribution of microperforate tenuitellid
1719 planktonic foraminifera in Holes 747A and 749B, Kerguelen Plateau. *Proceedings of*
1720 *the Ocean Drilling Program, Scientific Results* 120, 569-602.
- 1721 Li, Q., Jian, Z., Li, B., 2004. Oligocene-Miocene planktonic foraminifer biostratigraphy,
1722 Site 1148, northern South China Sea. In: Prell, W.L. et al. (Eds.), *Proceedings of the*
1723 *Ocean Drilling Program, Scientific Results* 184.
- 1724 Lourens, L.J., Hilgen, F.J., Shackleton, N.J., Laskar, J., Wilson, D., 2004. The Neogene
1725 Period. In: Gradstein, F.M., Ogg, J.G., Smith, A.G. (Eds.), *Geological Time Scale*

- 1726 2004. Cambridge University Press, pp. 409-440.
- 1727 Lowrie, W., Alvarez, W., Napoleone, G., Perch-Nielsen, K., Premoli Silva, I.,
1728 Toumarkine, M., 1982. Paleogene magnetic stratigraphy in Umbrian pelagic
1729 carbonate rocks: the Contessa sections, Gubbio. Geological Society of America
1730 Bulletin 93, 414–432.
- 1731 Luciani, V., Giusberti, L., Agnini, C., Fornaciari, E., Rio, D., Spofforth, D.J.A., Palike,
1732 H., 2010. Ecological and evolutionary response of Tethyan planktonic foraminifera
1733 to the middle Eocene climatic optimum (MECO) from the Alano section (NE Italy).
1734 Palaeogeography, Palaeoclimatology, Palaeoecology 292, 82-95.
- 1735 Luterbacher, H. P., Ali, J. R., Brinkhuis, H., Gradstein, F. M., Hooker, J., Monechi, S.,
1736 Ogg, J. G., Powell, J. Röhl, U., Sanfilippo, A., Schmitz, B., 2004. The Paleogene
1737 Period. In: Gradstein, F., Ogg, J., Smith, A., (Eds.), A Geologic Time Scale 2004:
1738 Cambridge University Press, 384–408.
- 1739 Lyle, M., Raffi, I., Pälike, H., Nishi, H., Gamage, K., Klaus, A., and the Expedition
1740 320/321 Scientists, 2009. Pacific Equatorial Transect IODP Preliminary Report, 321.
1741 doi:10.2204/iodp.pr.321.2009
- 1742 McGowran, B., 2005. Biostratigraphy: microfossils and geological time. Cambridge
1743 University Press, Cambridge, 459 pp.
- 1744 McGowran, B., Berggren, W.A., Hilgen, F., Steininger, F., Aubry, M.-P., Lourens, L.,
1745 Van Couvering, J., 2009. Neogene and Quaternary coexisting in the geological time
1746 scale: The inclusive compromise. Earth-Science Reviews 96, 249-262.
- 1747 Miller, K.G., Aubry, M.-P., Khan, M.J., Melillo, A.J., Kent, D.V., Berggren, W.A., 1985.
1748 Oligocene–Miocene biostratigraphy, magnetostratigraphy, and isotope stratigraphy

- 1749 of the western North Atlantic. *Geology* 13, 257-261.
- 1750 Miller, K.G., Feigenson, M.D., Wright, J.D., Clement, B.M., 1991. Miocene isotope
1751 reference section, Deep Sea Drilling Project Site 608: an evaluation of isotope and
1752 biostratigraphic resolution. *Paleoceanography*, 6: 33-52.
- 1753 Miller, K.G., Browning, J.V., Aubry, M-P., Wade, B.S., Katz, M.E., Kulpecz, A.A.,
1754 Wright, J.D., 2008. Eocene-Oligocene global climate and sea-level changes: St.
1755 Stephens Quarry, Alabama. *Geological Society America Bulletin* 120, 34-53.
- 1756 Mix, A.C., Le, J., Shackleton, N.J., 1995. Benthic foraminiferal stable isotope
1757 stratigraphy of Site 846: 0–1.8 Ma. In: Pisias, N.G., Mayer, L.A., Janecek, T.R.,
1758 Palmer-Julson, A., van Andel, T.H. (Eds.), *Proceeding of the Ocean Drilling*
1759 *Program, Scientific Results* 138, 839–854.
- 1760 Molina, E., Alegret, L., Apellaniz, E., Bernaola, G., Caballero, F., Dinares-Turell, J.,
1761 Hardenbol, J., Heilmann-Clausen, C., Larrasoña, J.C., Luterbacher, H., Monechi,
1762 S., Ortiz, S., Orue-Etxebarria, X., Payros, A., Pujalte, V., Rodríguez-Tovar, F.J.,
1763 Tori, F., Tosquella, J. and Uchman, A. 2010. Proposal for the Global Standard
1764 Stratotype-section and Point (GSSP) for the base of the Lutetian Stage at the
1765 Gorronatxe section (Spain). *Episodes*, in press.
- 1766 Muttoni, G., Kent, D.V., 2007. Widespread formation of cherts during the early Eocene
1767 climate optimum. *Palaeogeography, Palaeoclimatology, Palaeoecology* 253, 348–
1768 362.
- 1769 Nicholas, C.J., Pearson, P.N., Bown, P.R., Dunkley Jones T., Huber, B.T., Karega, A.,
1770 Lees, J.A., McMillan, I. K., O'Halloran, A., Singano, J.M., Wade, B.S., 2006.
1771 *Stratigraphy and sedimentology of the Upper Cretaceous to Paleogene Kilwa Group*,

- 1772 Tanzania. *Journal of African Earth Sciences* 45, 431-466.
- 1773 Olsson, R.K., 1972. Growth changes in the *Globorotalia foshi* lineage. *Eclogae*
1774 *Geologica Helvetica* 65 (1), 165–184.
- 1775 Olsson, R.K., Hemleben, C., Berggren, W.A., (Eds.), 1999. Atlas of Paleocene
1776 Planktonic Foraminifera. *Smithsonian Contributions to Paleobiology*, 85, 1–25.
- 1777 Ortiz, S., Gonzalvo, C., Molina, E., Rodríguez-Tovar, F.J., Uchman, A., Vandenberghe,
1778 N., Zeelmaekers, E., 2008, Palaeoenvironmental turnover across the Ypresian-
1779 Lutetian transition at the Agost section, southeastern Spain: in search of a marker
1780 event to define the Stratotype for the base of the Lutetian Stage. *Marine*
1781 *Micropaleontology* 69, 297-313.
- 1782 Ouda, K., Aubry, M.-P (Eds.), 2003. The Upper Paleocene-Lower Eocene of the Upper
1783 Nile Valley: Part 1: Stratigraphy: *Micropaleontology*, v. 49, Supplement 1, 212 p.
- 1784 Pälike, H., Norris, R.D., Herrle, J.O., Wilson, P.A., Coxall, H.K., Lear, C.H., Shackleton,
1785 N.J., Tripathi, A.K., Wade, B.S. 2006. The heartbeat of the Oligocene climate system.
1786 *Science* 314 (5807), 1894-1898.
- 1787 Pälike, H., Nishi, H., Lyle, M., Raffi, I., Klaus, A., Gamage, K., and the Expedition
1788 320/321 Scientists, 2009. Pacific Equatorial Transect. IODP Preliminary Report,
1789 320. doi:10.2204/iodp.pr.320.2009
- 1790 Payros, A., Bernaola, G., Orue-Etxebarria, X., Dinares-Turell, J., Tosquella, J., Apellaniz,
1791 E. 2007. Reassessment of the Early-Middle Eocene biomagnetochronology based on
1792 evidence from the Gorrondatxe section (Basque Country, western Pyrenees). *Lethaia*
1793 40, 183-195.
- 1794 Payros, A., Orue-Etxebarria, X., Bernaola, G., Apellaniz, E., Dinarès-Turell, J.,

- 1795 Tosquella, J. & Caballero, F. 2009. Characterization and astronomically calibrated
 1796 age of the first occurrence of *Turborotalia frontosa* in the Gorrondatxe section, a
 1797 prospective Lutetian GSSP: implications for the Eocene time scale. *Lethaia* 42, 255-
 1798 264.
- 1799 Pearson, P. N., 1995. Planktonic foraminifer biostratigraphy and the development of
 1800 pelagic caps on guyots in the Marshall Islands Group. In: Haggerty, J., Premoli
 1801 Silva, I., Rack, F., McNutt, M.K. (Eds.), *Proceeding of the Ocean Drilling Program,*
 1802 *Scientific Results* 144, 21-59.
- 1803 Pearson, P.N., 1998. Evolutionary concepts in biostratigraphy. In: Doyle, P., Bennett,
 1804 M.R., (Eds.), *Unlocking the Stratigraphical Record.* John Wiley & Sons, New York,
 1805 123-144
- 1806 Pearson, P. N., Chaisson, W. P. 1997. Late Paleocene to middle Miocene planktonic
 1807 foraminifer biostratigraphy of the Ceara Rise. In: Shackleton, N. J., Curry, W. B.,
 1808 Richter, C., Bralower, T. J. (Eds.), *Proceeding of the Ocean Drilling Program,*
 1809 *Scientific Results* 154, 33 - 68.
- 1810 Pearson, P.N., Wade, B.S., 2009. Taxonomy and stable isotope paleoecology of well-
 1811 preserved planktonic foraminifera from the uppermost Oligocene of Trinidad.
 1812 *Journal of Foraminifera Research* 39, 191-217.
- 1813 Pearson, P. N. Nicholas, C. J., Singano, J.M., Bown, P. R., Coxall, H.K., van Dongen, B.
 1814 E., Huber, B. T., Karega, A., Lees, J. A., Msaky, E., Pancost, R. D., Pearson, M.,
 1815 Roberts, A. P., 2004. Paleogene and Cretaceous sediment cores from the Kilwa and
 1816 Lindi areas of coastal Tanzania: Tanzania Drilling Project Sites 1 to 5. *Journal of*
 1817 *African Earth Sciences* 39, 25-62.

- 1818 Pearson, P.N., Olsson, R.K., Huber, B.T., Hemleben, C., Berggren, W.A. (Eds.), Atlas of
1819 Eocene Planktonic Foraminifera, Cushman Foundation Special Publication, 41.
- 1820 Poore, R. Z., Berggren, W.A., 1975. Late Cenozoic planktonic foraminiferal
1821 biostratigraphy and paleoclimatology of Hatton-Rockall Basin, DSDP Site 116, Jour.
1822 Foram. Research, 5, 270-293.
- 1823 Premoli Silva, I., Spezzaferri, S. 1990. Paleogene planktonic foraminifer biostratigraphy
1824 and paleoenvironmental remarks on Paleogene sediments from Indian Ocean Site,
1825 Leg 115. In: Duncan, R.A., Backman, J., Peterson, L.C., et al. Proceeding of the
1826 Ocean Drilling Program, Scientific Results 115, 277 - 314.
- 1827 Pujol, C., Duprat, J., 1983. Quaternary planktonic foraminifers of the southwestern
1828 Atlantic (Rio Grande Rise) Deep Sea Drilling Project Leg 72. In Barker, P.F.,
1829 Carlson, R.L., Johnson, D.A., et al., Initial Reports of the Deep Sea Drilling Project
1830 72, Washington (U.S. Govt. Printing Office), 601-615.
- 1831 Rögl, F., Egger, H. 2010. The missing link in the evolutionary origin of the foraminiferal
1832 genus *Hantkenina* and the problem of the lower-middle Eocene boundary. Geology
1833 38, 23-26.
- 1834 Shackleton, N. J., Crowhurst, S. 1997. Sediment fluxes based on an orbitally tuned time
1835 scale 5 Ma to 14 Ma, Site 926. In: Shackleton, N. J., Curry, W. B., Richter, C.,
1836 Bralower, T. J. (Eds.), Proceeding of the Ocean Drilling Program, Scientific Results
1837 154, 69 - 82.
- 1838 Shackleton, N.J., Berger, A., Peltier, W.R., 1990. An alternative astronomical calibration
1839 of the lower Pleistocene timescale based on ODP Site 677. Phil. Trans. R. Soc.
1840 Edinburgh, Earth Sciences 81, 251 – 261.

- 1841 Shackleton, N.J. Baldauf, J.G. Flores, J.-A. Iwai, M. Moore, Jr. T.C., Raffi, I., Vincent,
1842 E., 1995. Biostratigraphic summary for Leg 138. In: Pisias, N.G., Mayer, L.A.,
1843 Janacek, T.R., et al. (Eds.), Proceedings of the Ocean Drilling Program, Scientific
1844 Results, College Station, TX (Ocean Drilling Program), 138, pp. 517–536.
- 1845 Shackleton, N. J., Crowhurst, S. J., Weedon, G. P., Laskar, J. 1999. Astronomical
1846 calibration of Oligocene – Miocene time. *Philos. Trans. R. Soc. Lond. Ser. A: Math.*
1847 *Phys. Sci.* 357, 1907 – 1929.
- 1848 Shipboard Scientific Party, 1988. Sites 677 and 678. In: Becker, K., Sakai, H. et al.,
1849 (Eds.), Proceedings of the Ocean Drilling Program, Initial Reports 111, 253-346.
- 1850 Shipboard Scientific Party, 2002. Site 1218. In: Lyle, M., Wilson, P.A., Janacek, T.R., et
1851 al., Proceedings of the Ocean Drilling Program, Initial Reports, 199: College Station,
1852 TX (Ocean Drilling Program).
- 1853 Shipboard Scientific Party, 2004. Site 1264. In: Zachos, J.C., Kroon, D., Blum, P., et al.,
1854 Proceedings of the Ocean Drilling Program, Initial Reports, 208: College Station,
1855 TX (Ocean Drilling Program), 1–73.
- 1856 Shipboard Scientific Party, in press. In: Pälike, H., Lyle, M., Nishi, H., Raffi, I., Klaus,
1857 A., Gamage, K., et al., Proceedings of the Ocean Drilling Program, Initial Reports,
1858 320/321: College Station, TX (Ocean Drilling Program).
- 1859 Srinivasan, M.S., Chaturvedi, S.N., 1992. Neogene planktonic foraminiferal
1860 biochronology of the DSDP sites along the Ninetyeast Ridge, northern Indian
1861 Ocean. In: Ishizaki, K., Saito, T. (Eds.), Centenary of Japanese Micropaleontology,
1862 175-188.

- 1863 Srinivasan, M.S., Kennett, J.P., 1981a. A review of Neogene planktonic foraminiferal
1864 biostratigraphy: applications in the equatorial and South Pacific. In Warne, J.E.,
1865 Douglas, R.G., Winterer, E.L. (Eds.), *The Deep-Sea Drilling Project: A Decade of*
1866 *Progress. Spec. Publ.—Soc. Econ. Paleontol. Mineral.* 32, 395-432.
- 1867 Srinivasan, M.S., Kennett, J.P., 1981b. Neogene planktonic foraminiferal biostratigraphy
1868 and evolution: equatorial to subantarctic South Pacific. *Mar. Micropaleontol.* 6,
1869 499-533.
- 1870 Srinivasan, M.S., Sinha, D.K., 1992. Late Neogene planktonic foraminiferal events of the
1871 southwest Pacific and Indian Ocean: a comparison. In: Tsuchi, R., Ingle, Jr., J.C.
1872 (Eds.), *Pacific Neogene: Environment, Evolution and Events*: Tokyo (Univ. Tokyo
1873 Press), 203–220.
- 1874 Stainforth, R. M., Lamb, J. L., Luterbacher, H., Beard, J. H., Jeffords, R. M., 1975.
1875 *Cenozoic Planktonic Foraminiferal Zonation and Characteristics of Index Forms*:
1876 *University of Kansas Paleontological Contributions*, Article 62, 1-425.
- 1877 Steininger, F. F., Aubry, M. P., Berggren, W. A., Biolzi, M., Borsetti, A.M., Brzobohaty,
1878 R., Cartlidge, J.E., Cati, F., Corfield, R., Gelati, R., Iacarrino, S., Mödden, C.,
1879 Napoleone, D., Nolf, F., Ottner, B., Reichenbacher, B., Rögl, F., Roetzel, R.,
1880 Spezzaferri, S., Tateo, F., Villa, G., Wielandt, U., Zevenboom, D., Zorn, I., 1997.
1881 *The Global Stratotype Section and Point the GSSP for the Base of the Neogene*
1882 *(The Paleogene/Neogene Boundary)*. *Episodes* 20, 23-28.
- 1883 Stott, L. D., Kennett, J. P., 1990, Antarctic Paleogene planktonic foraminifer
1884 biostratigraphy: ODP Leg 113, Sites 689 and 690. In: Barker, P. F., Kennett, J. P.,

- 1885 et al. (Eds.), Proceedings of the Ocean Drilling Program 113, Ocean Drilling
 1886 Program, College Station, Texas, 549–569.
- 1887 Subbotina, N.N., 1953. Iskopaemye foraminifery SSSR (Globigerinidy, Khantkenininidy
 1888 i Globorotaliidy): Trudy Vsesoyznogo Nauchno-Issledovatel'skogo Geologo-
 1889 razvedochnogo Instituta (VNIGRI), 76, GOSTOPETEKHIZDAT, Leningrad, 295 p.
 1890 Translated into English by E. Lees, 1971, Fossil Foraminifera of the USSR:
 1891 Globigerinidae, Hantkeninidae and Globorotaliidae: Collet's Ltd., London and
 1892 Wellingborough, 321 p.
- 1893 Thompson, P.R., Bé, A.W.H., Duplessy, J.-C., Shackleton, N.J., 1979. Disappearance of
 1894 pink-pigmented *Globigerinoides ruber* at 120,000 yr BP in the Indian and Pacific
 1895 oceans. Nature 280, 554–558.
- 1896 Thunell, R., 1981. Late Miocene-early Pliocene planktonic foraminiferal biostratigraphy
 1897 and paleoceanography of low-latitude marine sequences. Marine Micropaleontology
 1898 6, 71-90.
- 1899 Turco, E., Bambini, A.M., Foresi, L.M., Iaccarino, S., Lirer, F., Mazzei, R., Salvatorini,
 1900 G., 2002, Middle Miocene high resolution calcareous plankton biostratigraphy at
 1901 Site 926 (Leg 154, equatorial Atlantic Ocean): paleoecological and
 1902 paleobiogeographical implications. Geobios 35, 257-276.
- 1903 Van Couvering, J.A., Castradori, D., Cita, M.B., Hilgen, F.J., Rio, D., 2000. The base of
 1904 the Zanclean Stage and of the Pliocene Series. Episodes 23, 179– 187.
- 1905 Van Couvering, J.A., Aubry, M.-P., Berggren, W.A., Gradstein, F.M., Hilgen, F.J., Kent,
 1906 D.V., Lourens, L.J., McGowran, B. What, if anything, is Quaternary. Stratigraphy
 1907 (in press).

- 1908 Wade, B. S., 2004. Planktonic foraminiferal biostratigraphy and mechanisms in the
1909 extinction of *Morozovella* in the late Middle Eocene. *Marine Micropaleontology* 51,
1910 23 – 38.
- 1911 Wade, B.S., Pälike, H., 2004. Oligocene climate dynamics. *Paleoceanography*, 19,
1912 PA4019, doi:10.1029/2004PA001042.
- 1913 Wade, B.S., Pearson, P.N., 2008. Planktonic foraminiferal turnover, diversity fluctuations
1914 and geochemical signals across the Eocene/Oligocene boundary in Tanzania. *Marine*
1915 *Micropaleontology* 68, 244-255.
- 1916 Wade, B.S., Olsson, R.K., 2009. Investigation of pre-extinction dwarfing in Cenozoic
1917 planktonic foraminifera. In: Wade, B.S. and Twitchett, R. (Eds.). *Extinction,*
1918 *Dwarfing and the Lilliput Effect*. *Palaeogeography, Palaeoclimatology,*
1919 *Palaeoecology*, Special Issue 284, 39-46.
- 1920 Wade, B.S., Berggren, W.A., Olsson, R.K. 2007. The biostratigraphy and paleobiology of
1921 Oligocene planktonic foraminifera from the equatorial Pacific Ocean (ODP Site
1922 1218). *Marine Micropaleontology* 62, 167-179.
- 1923 Weaver, P.P.E., Clement, B.M., 1987. Magnetobiostratigraphy of planktonic
1924 foraminiferal datums, DSDP Leg 94, North Atlantic. In: Ruddiman, W.F., Kidd,
1925 R.B., Thomas, E., et al., *Init. Repts. DSDP, 94: Washington (U.S. Govt. Printing*
1926 *Office)* 815–829.
- 1927 Zhang, J., Miller, K.G., Berggren, W.A., 1993. Neogene planktonic foraminiferal
1928 biostratigraphy of the northeastern Gulf of Mexico. *Micropaleontology* 39, 299-326.
- 1929 Zijdeveld, J.D.A., Hilgen, F.A., Langeres, C.G., Verhallen, P.J.J.M., Zachariasse, W.J.,
1930 1991. Integrated magnetostratigraphy and biostratigraphy of the upper Pliocene -

- 1931 lower Pleistocene from the Monte Singa and Crotone areas in Calabria, Italy. *Earth*
- 1932 and Planetary Sci. Letts. 107 697-714.

1933 **Table Caption**

1934 **Table 1.** Planktonic foraminiferal bioevents for the Cenozoic calibrated to the
1935 geomagnetic polarity time scale of Cande and Kent (1995). Marker taxa are highlighted
1936 in bold. Previously published ages are on multiple time scales and refer to the data in
1937 each given reference.

1938

1939 **Table 2.** Earth's ~405 kyr eccentricity cycle duration and ages for the Cenozoic with
1940 code names for the present to 41.5 Ma (base of Chron 19n).

1941

1942 **Table 3.** Planktonic foraminiferal bioevents for the Neogene Period calibrated to the
1943 geomagnetic polarity time scale and astronomical time scale of Lourens et al. (2004).

1944 Marker taxa are highlighted in bold.

1945

1946 **Table 4.** Planktonic foraminiferal bioevents for the Paleogene Period calibrated to the
1947 geomagnetic polarity time scale of Luterbacher et al. (2004) and astronomical time scale
1948 of Pälike et al. (2006). Marker taxa are highlighted in bold.

1949

1950

1951

1952

Figure Captions

Fig. 1. Nomenclature of biostratigraphical zones modified from Hedberg (1976) and Pearson (1998) to illustrate the convention of Berggren and Pearson (2005) and this paper. Note that examples C, D and E are described as 'interval zones' by the International Stratigraphic Guide (Hedberg, 1976) and all five examples are described as 'interval zones' by the North American Commission on Stratigraphy. We prefer to refer to the five different logical possibilities by different names. Note that our naming convention demands that the named species occur within the zone, hence the necessity of species C in example E.

Fig. 2. Primary planktonic foraminiferal bioevents for the Cenozoic against the polarity time scale of Cande and Kent (1995). BKSA95 = Berggren et al. (1995); BP05 = Berggren and Pearson (2005). (a) 0-25 Ma; (b) 20-45 Ma; (c) 40-65 Ma. A = Atlantic; IP = Indo-Pacific.

Fig. 3. Primary planktonic foraminiferal bioevents for the Neogene and late Paleogene against the astronomical time scale of Lourens et al. (2004, until base of Chron C6Cn.2n) and Pälike et al. (2006, from top Chron C6Cn.3n until base C19n). (a) 0-25 Ma; (b) 20-41.5 Ma. A = Atlantic; IP = Indo-Pacific. The ~405 kyr eccentricity cycle numbers are counted from the present.

Fig. 4. Primary planktonic foraminiferal bioevents for the Cenozoic against the polarity time scale of Gradstein et al. (2004). (a) 0-25 Ma; (b) 20-45 Ma; (c) 40-65 Ma. A =

1976 Atlantic; IP = Indo-Pacific.

1977

1978 **Fig. 5.** Revision of mid Miocene “M” Zones and comparison to previous zonations.

1979

1980 **Fig. 6.** Planktonic foraminiferal bioevents for the early-middle Eocene against the
1981 polarity time scale of Cande and Kent (1995). Primary marker taxa are in black,
1982 secondary markers and uncalibrated events to the magnetochronology are shown in grey.
1983 Planktonic foraminifera ranges are constrained from the Agost Section (Ortiz et al., 2008;
1984 Larrasoña et al., 2008) and Tanzania Drilling Project Site 2 (Pearson, unpublished data).

Zone (BKSA95 & BP05) (base)		Zone (This study) (base)		Datum		Notes		Age (Ma) This Study ^a		Published Age (Ma)		Reference		Berggren et al. (1995a,b)		Berggren & Pearson (2005)	
PT1b	PT1b	T	Globorotalia	flexuosa				0.07	0.07	Berggren	et	al.	1995 a	0.07	-		
		T	Globigerinoides	ruber (pink)	(Indo-Pacific only)			0.12	0.12	Thompson	et	al.	1979	-	-		
		B	Globigerinella	calida				0.22	0.22	Chaproniere	et	al.	1994	0.22	-		
		B	Globorotalia	flexuosa				0.40	0.40	Berggren	et	al.	1995 a	0.40	-		
		B	Globorotalia	hirsuta				0.45	0.45	Pujol	and	Duprat	1983	0.45	-		
		T	Globorotalia	tosaensis				0.61	0.61	Mix	et	al.	1995	0.65	-		
		B	Globorotalia	hessi				0.75	0.75	Chaproniere	et	al.	1994	0.75	-		
		X	random to dextral in Pulleniatina			(Pacific)		0.80	0.80	Pearson			1995	-	-		
		B	Globorotalia	excelsa				1.00	1.00	Berggren	et	al.	1995 a	1.00	-		
		T	Globoturborotalita	obliquus				1.30	1.30	Chaisson	and	Pearson	1997	-	-		
Pliocene/Pleistocene boundary																	
											1.80	Aguirre	and	Pasini	1985	-	-
PT1a	PT1a	T	Globigerinoides	fistulosus		k		1.88	1.88	Chaisson	and	Pearson	1997	1.77	-		
		B	Globorotalia	truncatulinoides		g		1.93	1.92	Chaisson	and	Pearson	1997	2.00	-		
		T	Globigerinoides	extremus				1.99	1.98	Chaisson	and	Pearson	1997	1.77	-		
		B	Pulleniatina	finalis				2.05	2.04	Chaisson	and	Pearson	1997	-	-		
		T	Globorotalia	exilis	(Atlantic)			2.10	2.09	Chaisson	and	Pearson	1997	2.15	-		
PL6	PL6 (Indo-Pacific)	Reappearance of Pulleniatina			(Atlantic)			2.26	2.26	Chaisson	and	Pearson	1997	2.30	-		
		T	Globoturborotalita	woodi				2.30	2.33	Chaisson	and	Pearson	1997	-	-		
		T	Globorotalia	pertenius				2.30	2.33	Chaisson	and	Pearson	1997	-	-		
		T	Globorotalia	pseudomiocenica	(Indo-Pacific)			2.30	2.30	Berggren	et	al.	1995 a	2.30	-		
		PL6	PL6 (Atlantic)	T	Globorotalia	miocenica	(Atlantic)		2.39	2.38	Chaisson	and	Pearson	1997	2.30	-	
		T	Globorotalia	limbata				2.39	2.38	Chaisson	and	Pearson	1997	-	-		
		T	Globorotalia	pertenuis				2.60	2.60	Berggren	et	al.	1995 a	2.60	-		
		T	Globoturborotalita	decoraperta				2.75	2.75	Chaisson	and	Pearson	1997	-	-		
		T	Globorotalia	multicamerata				2.99	3.10	Chaisson	and	Pearson	1997	3.09	-		
		PL5	PL5 (Atlantic)	T	Dentoglobigerina	altispira	(Atlantic) ^h		3.13	3.11	Chaisson	and	Pearson	1997	3.09	-	
PL4	PL4	T	Sphaeroidinellopsis	seminulina	(Atlantic) ^h		3.16	3.14	Chaisson	and	Pearson	1997	3.12	-			
		B	Globigerinoides	fistulosus				3.33	3.33	Berggren	et	al.	1995 a	3.33	-		
PL5	PL5 (Indo-Pacific)	B	Globorotalia	tosaensis				3.35	3.35	Berggren	et	al.	1995 a	3.35	-		
		T	Pulleniatina	disappearance	(Atlantic)			3.41	3.41	Chaisson	and	Pearson	1997	-	-		
		T	Dentoglobigerina	altispira	(Pacific)			3.46	3.05	Shackleton	et	al.	1995	-	-		
		B	Globorotalia	pertenuis				3.51	3.52	Chaisson	and	Pearson	1997	3.45	-		
		PL4	PL4	T	Sphaeroidinellopsis	seminulina	(Pacific)		3.57	3.20	Shackleton	et	al.	1995	-	-	
PL3	PL3	T	Pulleniatina	primalis				3.65	3.65	Berggren	et	al.	1995 a	3.65	-		
		B	Globorotalia	miocenica	(Atlantic)			3.76	3.77	Chaisson	and	Pearson	1997	3.55	-		
		T	Globorotalia	plesiotumida				3.76	3.77	Chaisson	and	Pearson	1997	-	-		
		T	Globorotalia	margaritae				3.84	3.85	Chaisson	and	Pearson	1997	3.58	-		
		X	Pulleniatina	sinis.-dext.				4.07	4.08	Chaisson	and	Pearson	1997	3.95	-		
		T	Pulleniatina	spectabilis	(Pacific)			4.20	4.20	Berggren	et	al.	1995 a	4.20	-		
		B	Globorotalia	crassaformis	sensu lato			4.30	4.31	Chaisson	and	Pearson	1997	4.50	-		

PL2	PL2	T	Globoturborotalita	nepenthes		4.36	4.39 Chaisson	and Pearson	1997	4.20	-
		B	Globorotalia	exilis		4.44	4.45 Chaisson	and Pearson	1997	-	-
		T	Sphaeroidinellopsis	kochi		4.52	4.53 Chaisson	and Pearson	1997	-	-
PL1b		T	Globorotalia	cibaoensis		4.60	4.60 Berggren	et al.	1995 b	4.60	-
<i>Miocene/Pliocene boundary</i>						5.33	5.33 <i>Van Couvering et al.</i>		2000	-	-
		B	Sphaeroidinella	dehiscens	sensu lato	5.48	5.54 Chaisson	and Pearson	1997	5.20	-
PL1a	PL1	B	Globorotalia	tumida	(Pacific)	5.51	5.59 Shackleton	et al.	1995	5.60	-
PL1a	PL1	B	Globorotalia	tumida	(Atlantic)	5.63	5.82 Chaisson	and Pearson	1997	5.60	-
		B	Turborotalita	humilis		5.71	5.84 Chaisson	and Pearson	1997	-	-
		T	Globoquadrina	dehiscens		5.80	5.80 Berggren	et al.	1995 a	5.80	-
		B	Globorotalia	margaritae		5.95	6.09 Chaisson	and Pearson	1997	6.40	-
M14	M14	T	Globorotalia	languaensis		6.00	6.00 Berggren	et al.	1995	6.00	-
		B	Globigerinoides	conglobatus		6.08	6.20 Chaisson	and Pearson	1997	-	-
		X	Neogloboquadrina	acostaensis	trans. sinis.-dext	6.20	6.20 Berggren	et al.	1995 b	6.20	-
		B	Pulleniatina	primalis		6.40	6.40 Berggren	et al.	1995 b	6.40	-
		X	Neogloboquadrina	acostaensis	trans. dext.-sinis	6.60	6.60 Berggren	et al.	1995 b	6.60	-
		X	Neogloboquadrina	atlantica	trans. dext.-sinis	6.80	6.80 Berggren	et al.	1995 b	6.80	-
		B	Neogloboquadrina	humerosa		8.50	8.50 Berggren	et al.	1995 b	8.50	-
M13b	M13b	B	Globorotalia	plesirotunda	c (1)	8.52	8.58 Chaisson	and Pearson	1997	-	-
		B	Globigerinoides	extremus	b	8.86	8.94 Turco	et al.	2002	-	-
		B	Globorotalia	cibaoensis		9.34	9.44 Chaisson	and Pearson	1997	-	-
		B	Globorotalia	juanai		9.62	9.76 Chaisson	and Pearson	1997	-	-
M13a	M13a	B	Neogloboquadrina	acostaensis		9.79	9.82 Chaisson	and Pearson	1997	10.90	-
M12	M12	T	Paragloborotalia	mayeri		10.53	10.49 Chaisson	and Pearson	1997	11.40	-
		B	Globorotalia	limbata		10.66	10.57 Chaisson	and Pearson	1997	-	-
		T	Cassigerinella	chipolensis	b	10.84	10.81 Turco	et al.	2002	-	-
		B	Globoturborotalita	apertura		11.12	11.19 Chaisson	and Pearson	1997	-	-
		B	Globoturborotalita	decoraperta		11.42	11.46 Chaisson	and Pearson	1997	-	-
		T	Globigerinoides	subquadratus	b	11.46	11.55 Turco	et al.	2002	-	-
M11	M11	B	Globoturborotalita	nepenthes	b	11.55	11.64 Turco	et al.	2002	11.80	-
M10	M10	T	Fohsella	fohsi		11.71	11.68 Chaisson	and Pearson	1997	11.90	-
		T	Globorotalia	praescitula		11.90	11.90 Berggren	et al.	1995	11.90	-
		B	Globorotalia	languaensis	b	12.89	12.39 Turco	et al.	2002	-	-
		B	Sphaeroidinellopsis	subdehiscens	b	13.00	13.03 Turco	et al.	2002	-	-
M9b	M9b	B	Fohsella	robusta		13.09	13.18 Chaisson	and Pearson	1997	12.30	-
M9a		B	Fohsella	lobata		-	12.50 Berggren	et al.	1995 b	12.50	-
		T	Cassigerinella	martinezpicoi	b	13.22	13.28 Turco	et al.	2002	-	-
M8	M9a	B	Fohsella	fohsi		13.34	13.42 Chaisson	and Pearson	1997	12.70	-
	M8	B	Fohsella	"praefohsi"		13.74	14.00 Pearson	and Chaisson	1997	-	-
		T	Fohsella	peripheroronda	b	13.77	13.87 Turco	et al.	2002	14.60	-
		T	Clavatorella	bermudezi		13.79	14.20 Pearson	and Chaisson	1997	-	-
		T	Globorotalia	archeomenardii	b	13.84	13.87 Turco	et al.	2002	-	-
M7	M7	B	Fohsella	peripheroacuta	b	14.23	14.02 Turco	et al.	2002	14.80	-
		B	Globorotalia	praemenardii		14.39	14.90 Pearson	and Chaisson	1997	-	-

Wade et al Table 1

M6	M6	T	Praeorbulina	sicana		14.56	14.40	Shackleton	et al.	1999	14.80	-
		T	Globigerinatella	insueta		14.69	15.00	Pearson	and Chaisson	1997	-	-
		B	Orbulina	suturalis		15.10	15.10	Berggren	et al.	1995 b	15.10	-
		B	Clavatorella	bermudezi	f	15.76	15.80	Pearson	and Chaisson	1997	-	-
M5b	M5b	B	Praeorbulina	circularis	e	16.00	16.00	Berggren	et al.	1995 b	16.00	-
		B	Globorotalia	archeomenardii		16.29	16.20	Pearson	and Chaisson	1997	-	-
		B	Praeorbulina	glomerosa	sensu stricto	16.29	16.20	Shackleton	et al.	1999	16.10	-
		B	Praeorbulina	curva		16.30	16.30	Berggren	et al.	1995 b	16.30	-
M5a	M5a	B	Praeorbulina	sicana		16.40	16.40	Berggren	et al.	1995 b	16.40	-
M4b	M4b	B	Fohsella	birnageae		16.70	16.70	Berggren	et al.	1995 b	16.70	-
M4a	M4a	B	Globorotalia	zealandica	j	17.30	17.30	Berggren	et al.	1995 b	17.30	-
		T	Catapsydrax	dissimilis		17.62	17.50	Shackleton	et al.	1999	17.30	-
		B	Globigerinatella	insueta s. str.		17.69	17.40	Pearson	and Chaisson	1997	(18.8)	-
		B	Globorotalia	praescitula		18.50	18.50	Berggren	et al.	1995 b	18.50	-
M3	M3	T	Globoquadrina	binaiensis		19.43	19.10	Pearson	and Chaisson	1997	-	-
		B	Globigerinatella	sp.		19.66	20.20	Pearson	and Chaisson	1997	-	-
		B	Globigerinoides	altiaperturus		20.50	20.50	Berggren	et al.	1995 b	20.50	-
		T	Tenuitella	munda		21.40	21.40	Berggren	et al.	1995	21.40	-
M2	M2	T	Globigerina	angulisuturalis		21.60	21.60	Berggren	et al.	1995 b	21.60	-
		T	'Paragloborotalia'	kugleri		21.81	21.00	Shackleton	et al.	1999	21.50	-
		T	'Paragloborotalia'	pseudokugleri		22.04	21.10	Shackleton	et al.	1999	21.60	-
		B	Globoquadrina	dehiscens		23.20	23.20	Berggren	et al.	1995 b	23.20	-
M1b	M1b	T	Globigerina	ciperoensis		23.68	22.80	Shackleton	et al.	1999	-	-
		B	Globigerinoides	trilobus	sensu lato	23.73	22.90	Shackleton	et al.	1999	-	-
		B	'Paragloborotalia'	kugleri		23.73	22.90	Shackleton	et al.	1999	23.80	-
		Oligocene/Miocene boundary						23.80	Berggren	et al.	1995 b	
O7	O7	T	Tenuitella	gemma		24.3	24.3	Berggren	et al.	1995 b	24.3	-
		LCO	Globigerinoides	primordius		24.3	24.3	Berggren	et al.	1995 b	24.3	-
		B	'Paragloborotalia'	pseudokugleri		25.9	25.9	Berggren	et al.	1995 b	25.9	-
		B	Globigerinoides	primordius		26.7	26.7	Berggren	et al.	1995 b	26.7	-
O6	O6	T	Paragloborotalia	opima		27.5	27.5	Wade	et al.	2007	27.1	27.1
O5	O5	HCO	Chiloguembelina	cubensis		28.4	28.4	Wade	et al.	2007	28.5	28.5
O4	O4	B	Globigerina	angulisuturalis		29.4	29.4	Berggren	et al.	1995 b	29.4	29.4
O3	O3	T	Subbotina	angiporoides		30.0	30.0	Berggren	et al.	1995 b	30.0	30.0
		T	Turborotalia	ampliapertura		30.3	30.3	Berggren	et al.	1995 b	30.3	30.3
		B	Paragloborotalia	opima		30.6	30.6	Berggren	et al.	1995 b	30.6	-
		T	Pseudohastigerina	nagewichiensis		32.0	32.0	Berggren	et al.	1995 b	32.0	32.0
Eocene/Oligocene boundary												
O1	O1	T	Hantkenina	alabamensis		33.7	33.7	Berggren	and Pearson	2005	33.7	33.7
E16	E16	HCO	Pseudohastigerina	micra		33.7	33.7	Wade	and Pearson	2008	-	-
		T	Turborotalia	cerroazulensis		33.8	33.8	Berggren	and Pearson	2005	33.8	33.8
		T	Globigerinatheka	index		34.3	34.3	Berggren	and Pearson	2005	34.3	34.3
		B	Turborotalia	cunialensis		35.3	35.3	Berggren	and Pearson	2005	35.2	35.3
E15	E15	T	Globigerinatheka	semiinvoluta		35.8	35.8	Berggren	and Pearson	2005	35.3	35.8
		B	Globigerinatheka	semiinvoluta		38.0	38.0	Wade		2004	38.4	-

E14	E14	T	Morozovelloides	crassatus	38.0	38.0	Wade		2004	38.1	38.0
		T	Acarinina	mcgowrani	38.0	38.0	Wade		2004	-	-
		T	Turborotalia	frontosa	39.3	39.3	Berggren	et al.	1995 b	39.3	-
E13	E13	T	Orbulinoides	beckmanni	40.0	40.0	Wade		2004	40.1	40.0
E12	E12	B	Orbulinoides	beckmanni	40.5	40.5	Berggren	et al.	1995 b	40.5	40.5
		T	Acarinina	bullbrookii	40.5	40.5	Berggren	et al.	1995 b	40.5	-
E11	E11	T	Guembilitrioides	nuttalli	(42.3)	(42.3)	Berggren	and Pearson	2005	-	(42.3)
		B	Turborotalia	pomeroli	42.4	42.4	Berggren	et al.	1995 b	42.4	-
		B	Globigerinatheka	index	42.9	42.9	Berggren	et al.	1995 b	42.9	-
		B	Morozovelloides	lehneri	43.5	43.5	Berggren	et al.	1995 b	43.5	-
E10	E10	T	Morozovella	aragonensis	43.6	43.6	Berggren	et al.	1995 b	43.6	43.6
E9	E9	B	Globigerinatheka	kugleri	(44.4)	-	Pearson	et al.	2004	45.8	45.8
		B	Hantkenina	singanoae	44.5	43.5	Payros	et al.	2009	49.0	-
E8	E8	B	Guembilitrioides	nuttalli	46.4	45.5	Payros	et al.	2009	-	49.0
	E7b	B	Turborotalia	frontosa	49.0	48.6	Payros	et al.	2009	-	-
E7	E7a	B	Acarinina	cuneicamerata	50.4	50.4	Hancock	et al.	2002	-	50.4
		B	Planorotalites	palmerae	50.4	50.4	Berggren	et al.	1995 b	50.4	-
E6	E6	T	Morozovella	subbotinae	50.8	50.8	Berggren	and Pearson	2005	-	50.8
E5	E5	B	Morozovella	aragonensis	52.3	52.3	Berggren	et al.	1995 b	52.3	52.3
		T	Morozovella	marginodentata	52.5	52.5	Berggren	et al.	1995 b	52.5	-
		T	Morozovella	lensiformis	52.7	52.7	Berggren	et al.	1995 b	52.7	-
		T	Morozovella	aequa	53.6	53.6	Berggren	et al.	1995 b	53.6	-
E4	E4	B	Morozovella	formosa	54.0	54.0	Berggren	et al.	1995 b	54.0	54.0
		B	Morozovella	lensiformis	54.0	54.0	Berggren	et al.	1995 b	54.0	-
E3	E3	T	Morozovella	velascoensis	54.5	54.5	Berggren	and Pearson	2005	54.7	54.5
		T	Morozovella	acuta	54.7	54.7	Berggren	et al.	1995 b	54.7	-
		B	Morozovella	gracilis	54.7	54.7	Berggren	et al.	1995 b	54.7	-
		B	Igorina	broedermanni	54.7	54.7	Berggren	et al.	1995 b	54.7	-
		B	Morozovella	marginodentata	54.8	54.8	Berggren	et al.	1995 b	54.8	-
E2	E2	B	Pseudohastigerina	wilcoxensis	55.4	55.4	Berggren	and Pearson	2005	-	55.4
		B	Globanomalina	australiformis	55.5	55.5	Berggren	et al.	1995 b	55.5	-
E1	E1	B	Acarinina	sibaiyaensis	55.5	55.5	Berggren	and Pearson	2005	-	55.5
<i>Paleocene/Eocene boundary</i>						55.5	Ouda	and Aubry	2003		
P5	P5	T	Globanomalina	pseudomenardii	55.9	55.9	Berggren	et al.	1995 b	55.9	55.9
		B	Morozovella	subbotinae	55.9	55.9	Berggren	et al.	1995 b	55.9	-
		T	Acarinina	mckannai	56.3	56.3	Berggren	et al.	1995 b	56.3	-
		T	Acarinina	acarinata	56.3	56.3	Berggren	et al.	1995 b	56.3	-
P4c	P4c	B	Acarinina	soldadoensis	56.5	56.5	Berggren	et al.	1995 b	56.5	56.5
		B	Acarinina	coalingensis	56.5	56.5	Berggren	et al.	1995 b	56.5	-
		B	Morozovella	aequa	56.5	56.5	Berggren	et al.	1995 b	56.5	-
		T	Acarinina	subsphaerica	57.1	57.1	Berggren	et al.	1995 b	57.1	57.1
		B	Acarinina	mckannai	59.1	59.1	Berggren	et al.	1995 b	59.1	-
P4b	P4b	T	Parasubbotina	variospira	59.2	59.2	Berggren	et al.	1995 b	59.2	59.2
		B	Acarinina	acarinata	59.2	59.2	Berggren	et al.	1995 b	59.2	-
		B	Acarinina	subsphaerica	59.2	59.2	Berggren	et al.	1995 b	59.2	-

P4a	P4a	B	Globanomalina	pseudomenardii	59.4	59.4	Berggren	et al.	2000	59.2	59.4
P3b	P3b	B	Igorina	albeari	60.0	60.0	Berggren	et al.	1995 b	60.0	60.0
		B	Morozovella	velascoensis	60.0	60.0	Berggren	et al.	1995 b	60.0	-
		B	Morozovella	conicotruncata	60.9	60.9	Berggren	et al.	1995 b	60.9	-
P3a	P3a	B	Morozovella	angulata	61.0	61.0	Berggren	et al.	1995 b	61.0	61.0
		B	Igorina	pusilla	61.0	61.0	Berggren	et al.	1995 b	61.0	-
		B	Morozovella	praeangulata	61.2	61.2	Berggren	et al.	1995 b	61.2	-
		B	Globanomalina	imitata	(61.3)	(61.3)	Berggren	et al.	1995 b	(61.3)	-
P2	P2	B	Praemurica	uncinata	61.4	61.4	Berggren	and Pearson	2005	61.2	61.4
P1c	P1c	B	Globanomalina	compressa	62.9	62.9	Berggren	and Pearson	2005	63.0	62.9
		B	Praemurica	inconstans	62.9	62.9	Berggren	and Pearson	2005	63.0	62.9
		B	Parasubbotina	variata	63.0	63.0	Berggren	et al.	1995 b	63.0	-
P1b	P1b	B	Subbotina	triloculinoides	64.3	64.3	Berggren	et al.	1995 b	64.3	64.3
P1a	P1a	T	Parvularugoglobigerina	eugubina	64.8	64.8	Berggren	and Pearson	2005	64.7	64.8
		B	Parvularugoglobigerina	extensa	64.9	64.9	Olsson	et al.	1999	-	-
Pα	Pα	B	Parvularugoglobigerina	eugubina	64.97	64.97	Berggren	et al.	1995 b	64.97	64.97
P0	P0	T	Globotruncana		65.0	65.0	Berggren	et al.	1995 b	65.0	65.0

Notes

a	Calibrated to Cande and Kent (1995)
b	Calibration of Turco et al. (2002) on Astronomical Time Scale of Shackleton and Crowhurst (1997)
c	Replaces the and/or non zone of (1) BKSA95; (2) Berggren & Pearson (2005)
d	Note mistake in Berggren & Pearson (2005)
e	Note mistake in Lourens et al. (2004) Table A2.3, where Bottom <i>Praeorbulina circularis</i> should read Top <i>Praeorbulina circularis</i>
f	Note mistake in Shackleton et al. (1999) where 14.8 Ma should read 15.8 Ma. This mistake is also propagated in Lourens et al. (2004)
g	Traditionally used as base N22, this datum is considered highly diachronous between ocean basins (see Dowsett, 1988).
h	Age adjusted to reflect the short stratigraphic duration between <i>D. altispira</i> and <i>S. seminulina</i> - see text for discussion
i	Calibrated to Gradstein et al. (2004)
j	Note mistake in BKSA95 (page 174) where LAD should read FAD
k	Note mistake in Berggren et al. (1995a, Table 6) where 1.6 Ma should read 1.77 Ma
X	Change in coiling direction
HCO	Highest common occurrence
LCO	Lowest common occurrence
(44.4)	Estimated age

~405 kyr Cycle cycle Code/Chron	Younger end of cycle (Ma)	Older end of cycle (Ma)	Duration	Notes
1	-0.121	0.297	0.418	a
2 Pt-C1n	0.297	0.702	0.405	
3 Pt-C1n	0.702	1.106	0.405	
4 Pt-C1r	1.106	1.516	0.410	
5 Pt-C1r	1.516	1.922	0.406	
6 Pl-C2n	1.922	2.329	0.408	
7 Pl-C2r	2.329	2.736	0.407	
8 Pl-C2An	2.736	3.144	0.408	
9 Pl-C2An	3.144	3.543	0.399	
10 Pl-C2An	3.543	3.944	0.401	
11 Pl-C2Ar	3.944	4.341	0.397	
12 Pl-C3n	4.341	4.742	0.402	
13 Pl-C3n	4.742	5.155	0.413	
14 Pl-C3n	5.155	5.559	0.404	
15 Mi-C3r	5.559	5.968	0.409	
16 Mi-C3r	5.968	6.374	0.407	
17 Mi-C3An	6.374	6.765	0.391	
18 Mi-C3Ar	6.765	7.172	0.406	
19 Mi-C3Bn	7.172	7.574	0.403	
20 Mi-C4n	7.574	7.983	0.409	
21 Mi-C4n	7.983	8.395	0.412	
22 Mi-C4r	8.395	8.797	0.402	
23 Mi-C4An	8.797	9.205	0.409	
24 Mi-C4Ar	9.205	9.602	0.397	
25 Mi-C4Ar	9.602	10.008	0.406	
26 Mi-C5n	10.008	10.423	0.415	
27 Mi-C5n	10.423	10.834	0.412	
28 Mi-C5n	10.834	11.240	0.406	
29 Mi-C5r	11.240	11.644	0.405	
30 Mi-C5r	11.644	12.059	0.414	
31 Mi-C5An	12.059	12.461	0.403	
32 Mi-C5Ar	12.461	12.870	0.409	
33 Mi-C5Ar	12.870	13.273	0.403	
34 Mi-C5AAr	13.273	13.678	0.405	
35 Mi-C5ABr	13.678	14.078	0.400	
36 Mi-C5ACn	14.078	14.479	0.401	
37 Mi-C5ADn	14.479	14.895	0.416	
38 Mi-C5Bn	14.895	15.297	0.403	
39 Mi-C5Br	15.297	15.701	0.403	
40 Mi-C5Br	15.701	16.099	0.399	
41 Mi-C5Cn	16.099	16.493	0.393	
42 Mi-C5Cn	16.493	16.904	0.411	
43 Mi-C5Cr	16.904	17.308	0.404	
44 Mi-C5Dn	17.308	17.718	0.410	
45 Mi-C5Dr	17.718	18.134	0.416	
46 Mi-C5En	18.134	18.528	0.393	
47 Mi-C5Er	18.528	18.930	0.402	

Wade et al Table 2

48 Mi-C6n	18.930	19.328	0.399
49 Mi-C6n	19.328	19.736	0.408
50 Mi-C6r	19.736	20.150	0.414
51 Mi-C6An	20.150	20.559	0.409
52 Mi-C6An	20.559	20.965	0.406
53 Mi-C6Ar	20.965	21.366	0.401
54 Mi-C6AAr	21.366	21.778	0.412
55 Mi-C6Bn	21.778	22.183	0.405
56 Mi-C6Bn	22.183	22.595	0.412
57 Mi-C6Cn	22.595	22.996	0.401
58 Ol-C6Cn	22.996	23.401	0.406
59 Ol-C6Cn	23.401	23.809	0.407
60 Ol-C6Cr	23.809	24.210	0.401
61 Ol-C7n	24.210	24.623	0.414
62 Ol-C7r	24.623	25.031	0.408
63 Ol-C7Ar	25.031	25.435	0.404
64 Ol-C8n	25.435	25.824	0.389
65 Ol-C8n	25.824	26.221	0.397
66 Ol-C8r	26.221	26.633	0.412
67 Ol-C9n	26.633	27.041	0.408
68 Ol-C9n	27.041	27.452	0.411
69 Ol-C9r	27.452	27.856	0.405
70 Ol-C10n	27.856	28.257	0.400
71 Ol-C10r	28.257	28.657	0.400
72 Ol-C10r	28.657	29.048	0.392
73 Ol-C10r	29.048	29.454	0.405
74 Ol-C11n	29.454	29.861	0.407
75 Ol-C11n	29.861	30.274	0.413
76 Ol-C11r	30.274	30.681	0.407
77 Ol-C12n	30.681	31.081	0.400
78 Ol-C12r	31.081	31.488	0.407
79 Ol-C12r	31.488	31.892	0.404
80 Ol-C12r	31.892	32.303	0.411
81 Ol-C12r	32.303	32.708	0.404
82 Ol-C12r	32.708	33.120	0.413
83 Ol-C12r	33.120	33.523	0.403
84 Ol-C13n	33.523	33.921	0.398
85 Eo-C13r	33.921	34.322	0.401
86 Eo-C13r	34.322	34.734	0.412
87 Eo-C13r	34.734	35.151	0.417
88 Eo-C15n	35.151	35.548	0.398
89 Eo-C16n	35.548	35.956	0.407
90 Eo-C16n	35.956	36.351	0.395
91 Eo-C16n	36.351	36.751	0.401
92 Eo-C17n	36.751	37.158	0.406
93 Eo-C17n	37.158	37.557	0.399
94 Eo-C17n	37.557	37.971	0.414
95 Eo-C17n	37.971	38.369	0.398
96 Eo-C17r	38.369	38.768	0.399

Wade et al Table 2

97 Eo-C18n	38.768	39.179	0.410	
98 Eo-C18n	39.179	39.576	0.397	
99 Eo-C18n	39.576	39.979	0.403	
100 Eo-C18r	39.979	40.374	0.395	
101 Eo-C18r	40.374	40.781	0.407	
102 Eo-C18r	40.781	41.193	0.412	
103 Eo-C18r	41.193	41.601	0.408	
104	41.601	42.008	0.407	b
105	42.008	42.407	0.400	
106	42.407	42.814	0.407	
107	42.814	43.213	0.399	
108	43.213	43.625	0.413	
109	43.625	44.034	0.409	
110	44.034	44.442	0.408	
111	44.442	44.841	0.399	
112	44.841	45.238	0.396	
113	45.238	45.648	0.410	
114	45.648	46.063	0.415	
115	46.063	46.477	0.415	
116	46.477	46.875	0.398	
117	46.875	47.278	0.403	
118	47.278	47.675	0.397	
119	47.675	48.077	0.402	
120	48.077	48.485	0.409	
121	48.485	48.886	0.401	
122	48.886	49.297	0.411	
123	49.297	49.695	0.398	
124	49.695	50.100	0.405	
125	50.100	50.508	0.408	
126	50.508	50.899	0.391	
127	50.899	51.303	0.404	
128	51.303	51.704	0.401	
129	51.704	52.113	0.409	
130	52.113	52.532	0.419	
131	52.532	52.930	0.398	
132	52.930	53.331	0.401	
133	53.331	53.724	0.394	
134	53.724	54.129	0.405	
135	54.129	54.541	0.412	
136	54.541	54.953	0.412	
137	54.953	55.364	0.412	
138	55.364	55.763	0.399	
139	55.763	56.166	0.402	
140	56.166	56.568	0.402	
141	56.568	56.982	0.414	
142	56.982	57.388	0.406	
143	57.388	57.792	0.404	
144	57.792	58.202	0.410	
145	58.202	58.608	0.407	

Wade et al Table 2

146	58.608	59.020	0.412
147	59.020	59.419	0.399
148	59.419	59.824	0.404
149	59.824	60.231	0.408
150	60.231	60.639	0.407
151	60.639	61.055	0.417
152	61.055	61.462	0.407
153	61.462	61.868	0.406
154	61.868	62.261	0.392
155	62.261	62.664	0.404
156	62.664	63.072	0.407
157	63.072	63.482	0.410
158	63.482	63.896	0.413
159	63.896	64.291	0.395
160	64.291	64.693	0.401
161	64.693	65.089	0.396
162	65.089	65.494	0.405
163	65.494	65.910	0.416
164	65.910	66.313	0.403
165	66.313	66.724	0.411
166	66.724	67.121	0.397
167	67.121	67.520	0.398

Notes

^a This cycle ends in the future

^b Cycles below Chron C19n are not named pending development on the Paleogene astronomical time scale.

Zone (BKSA 95 & BP05) (base)	Zone (This study) (base)	Datum	Notes	Astronomical Age ^a	Reference	Published Age (Ma)	Reference	Berggren et al. (1995a,b)
		T	Globorotalia flexuosa	0.07	This study	0.07	Berggren et al. 1995 a	0.07
		T	Globigerinoides ruber (pink)	0.12	This study	0.12	Thompson et al. 1979	-
		B	Globigerinella calida	0.22	This study	0.22	Chaproniere et al. 1994	0.22
		B	Globorotalia flexuosa	0.40	This study	0.40	Berggren et al. 1995 a	0.40
		B	Globorotalia hirsuta	0.45	This study	0.45	Pujol and Duprat 1983	0.45
PT1b	PT1b	T	Globorotalia tosaensis	0.61	Lourens et al. 2004	0.61	Mix et al. 1995	0.65
		B	Globorotalia hessi	0.75	This study	0.75	Chaproniere et al. 1994	0.75
		X	random to dextral in Pulleniatina	0.80	This study	0.80	Pearson 1995	-
		B	Globorotalia excelsa	1.00	This study	1.00	Berggren et al. 1995 a	1.00
		T	Globoturborotalita obliquus	1.30	Lourens et al. 2004	1.30	Chaisson and Pearson 1997	-
		T	Globoturborotalita apertura	1.64	Lourens et al. 2004	1.64	Chaisson and Pearson 1997	-
<i>Pliocene/Pleistocene boundary</i>								
						1.81	Aguirre and Pasini 1985	-
PT1a	PT1a	T	Globigerinoides fistulosus	1.88	Lourens et al. 2004	1.88	Chaisson and Pearson 1997	1.77
		B	Globorotalia truncatulinoides	1.93	Lourens et al. 2004	1.92	Chaisson and Pearson 1997	2.00
		T	Globigerinoides extremus	1.98	Lourens et al. 2004	1.98	Chaisson and Pearson 1997	1.77
		B	Pulleniatina finalis	2.04	Lourens et al. 2004	2.04	Chaisson and Pearson 1997	-
		T	Globorotalia exilis	2.09	Lourens et al. 2004	2.09	Chaisson and Pearson 1997	2.15
			Reappearance of Pulleniatina	2.26	Lourens et al. 2004	2.26	Chaisson and Pearson 1997	2.30
		T	Globoturborotalita woodi	2.30	Lourens et al. 2004	2.33	Chaisson and Pearson 1997	-
		T	Globorotalia pertenuis	2.30	Lourens et al. 2004	2.33	Chaisson and Pearson 1997	-
PL6	PL6 (Indo-Pacific)	T	Globorotalia pseudomiocenica	2.30	This study	2.30	Berggren et al. 1995 a	2.30
PL6	PL6 (Atlantic)	T	Globorotalia miocenica	2.39	Lourens et al. 2004	2.38	Chaisson and Pearson 1997	2.30
		T	Globorotalia limbata	2.39	Lourens et al. 2004	2.38	Chaisson and Pearson 1997	-
		T	Globorotalia pertenuis	2.60	This study	2.60	Berggren et al. 1995 a	2.60
		T	Globoturborotalita decoraperta	2.75	Lourens et al. 2004	2.75	Chaisson and Pearson 1997	-
		T	Globorotalia multicamerata	2.98	Lourens et al. 2004	3.10	Chaisson and Pearson 1997	3.09
PL5	PL5 (Atlantic)	T	Dentoglobigerina altispira	3.13	Lourens et al. 2004	3.11	Chaisson and Pearson 1997	3.09
PL4	PL4	T	Sphaeroidinellopsis seminulina	3.16	Lourens et al. 2004	3.14	Chaisson and Pearson 1997	3.12
		B	Globigerinoides fistulosus	3.33	This study	3.33	Berggren et al. 1995 a	3.33
		B	Globorotalia tosaensis	3.35	This study	3.35	Berggren et al. 1995 a	3.35
		T	Pulleniatina disappearance	3.41	Lourens et al. 2004	3.41	Chaisson and Pearson 1997	-
PL5 (Indo-Pacific)	PL5 (Indo-Pacific)	T	Dentoglobigerina altispira	3.47	Lourens et al. 2004	3.05	Shackleton et al. 1995	-
		B	Globorotalia pertenuis	3.52	Lourens et al. 2004	3.52	Chaisson and Pearson 1997	3.45
PL4	PL4	T	Sphaeroidinellopsis seminulina	3.59	Lourens et al. 2004	3.20	Shackleton et al. 1995	-
		T	Pulleniatina primalis	3.66	This study	3.65	Berggren et al. 1995 a	3.65
		B	Globorotalia miocenica	3.77	Lourens et al. 2004	3.77	Chaisson and Pearson 1997	3.55
		T	Globorotalia plesiotumida	3.77	Lourens et al. 2004	3.77	Chaisson and Pearson 1997	-
PL3	PL3	T	Globorotalia margaritae	3.85	Lourens et al. 2004	3.85	Chaisson and Pearson 1997	3.58

Wade et al Table 3

		X	Pulleniatina	sinis.-dext.		4.08	Lourens	et al.	2004	4.08	Chaisson	and Pearson	1997	3.95
		T	Pulleniatina	spectabilis	(Pacific)	4.21	This study			4.20	Berggren	et al.	1995 a	4.20
		B	Globorotalia	crassaformis	sensu lato	4.31	Lourens	et al.	2004	4.31	Chaisson	and Pearson	1997	4.50
PL2	PL2	T	Globoturborotalita	nepenthes		4.37	Lourens	et al.	2004	4.39	Chaisson	and Pearson	1997	4.20
		B	Globorotalia	exilis		4.45	Lourens	et al.	2004	4.45	Chaisson	and Pearson	1997	-
		T	Sphaeroidinellopsis	kochi		4.53	Lourens	et al.	2004	4.53	Chaisson	and Pearson	1997	-
PL1b		T	Globorotalia	cibaoensis		4.61	This study			4.60	Berggren	et al.	1995 b	4.60
<i>Miocene/Pliocene boundary</i>														
		B	Sphaeroidinella	dehiscens	sensu lato	5.33	Lourens	et al.	2004	5.33	<i>Van Couvervi</i>	<i>et al.</i>	2000	-
PL1a	PL1	B	Globorotalia	tumida	(Pacific)	5.53	Lourens	et al.	2004	5.54	Chaisson	and Pearson	1997	5.20
PL1a	PL1	B	Globorotalia	tumida	(Atlantic)	5.57	Lourens	et al.	2004	5.59	Shackleton	et al.	1995	5.60
		B	Turborotalita	humilis		5.72	Lourens	et al.	2004	5.82	Chaisson	and Pearson	1997	5.60
		B	Globoquadrina	dehiscens		5.81	Lourens	et al.	2004	5.84	Chaisson	and Pearson	1997	-
		B	Globorotalia	margaritae		5.92	This study			5.80	Berggren	et al.	1995 a	5.80
M14	M14	T	Globorotalia	lenguaensis		6.08	Lourens	et al.	2004	6.09	Chaisson	and Pearson	1997	6.40
		B	Globigerinoides	conglobatus		6.13	This study			6.00	Berggren	et al.	1995	6.00
		X	Neogloboquadrina	acostaensis	trans. sinis.-dext.	6.20	Lourens	et al.	2004	6.20	Chaisson	and Pearson	1997	-
		B	Pulleniatina	primalis		6.34	This study			6.20	Berggren	et al.	1995 b	6.20
		X	Neogloboquadrina	acostaensis	trans. dext.-sinis.	6.60	This study			6.40	Berggren	et al.	1995 b	6.40
		X	Neogloboquadrina	atlantica	trans. dext.-sinis.	6.77	This study			6.60	Berggren	et al.	1995 b	6.60
		B	Neogloboquadrina	humerosa		6.99	This study			6.80	Berggren	et al.	1995 b	6.80
M13b	M13b	B	Globorotalia	plesiotumida	^c	8.56	This study			8.50	Berggren	et al.	1995 b	8.50
		B	Globigerinoides	extremus	^b	8.58	Lourens	et al.	2004	8.58	Chaisson	and Pearson	1997	-
		B	Globorotalia	cibaoensis		8.93	Lourens	et al.	2004	8.94	Turco	et al.	2002	-
		B	Globorotalia	juanai		9.44	Lourens	et al.	2004	9.44	Chaisson	and Pearson	1997	-
M13a	M13a	B	Neogloboquadrina	acostaensis		9.69	Lourens	et al.	2004	9.76	Chaisson	and Pearson	1997	-
M12	M12	T	Paragloborotalia	mayeri		9.83	Lourens	et al.	2004	9.82	Chaisson	and Pearson	1997	10.90
		B	Globorotalia	limbata		10.46	Lourens	et al.	2004	10.49	Chaisson	and Pearson	1997	11.40
		T	Cassigerinella	chipolensis	^b	10.64	Lourens	et al.	2004	10.57	Chaisson	and Pearson	1997	-
		B	Globoturborotalita	apertura		10.89	Lourens	et al.	2004	10.81	Turco	et al.	2002	-
		B	Globoturborotalita	decoraperta		11.18	Lourens	et al.	2004	11.19	Chaisson	and Pearson	1997	-
		T	Globigerinoides	subquadratus	^b	11.49	Lourens	et al.	2004	11.46	Chaisson	and Pearson	1997	-
M11	M11	B	Globoturborotalita	nepenthes	^b	11.54	Lourens	et al.	2004	11.55	Turco	et al.	2002	-
M10	M10	T	Fohsella	fohsi		11.63	Lourens	et al.	2004	11.64	Turco	et al.	2002	11.80
		T	Globorotalia	praescitula		11.79	Lourens	et al.	2004	11.68	Chaisson	and Pearson	1997	11.90
		B	Globorotalia	lenguaensis	^b	13.73	Lourens	et al.	2004	11.90	Berggren	et al.	1995	11.90
		B	Sphaeroidinellopsis	subdehiscens	^b	12.84	Lourens	et al.	2004	12.39	Turco	et al.	2002	-
M9b	M9b	B	Fohsella	robusta		13.02	Lourens	et al.	2004	13.03	Turco	et al.	2002	-
M9a		B	Fohsella	lobata		13.13	Lourens	et al.	2004	13.18	Chaisson	and Pearson	1997	12.30
		T	Cassigerinella	martinezpicoi	^b	-	-	-	-	12.50	Berggren	et al.	1995 b	12.50
M8	M9a	B	Fohsella	fohsi		13.27	Lourens	et al.	2004	13.28	Turco	et al.	2002	-
	M8	B	Fohsella	"praefohsi"		13.41	Lourens	et al.	2004	13.42	Chaisson	and Pearson	1997	12.70
		T	Fohsella	peripheroronda	^b	13.77	Lourens	et al.	2004	14.00	Pearson	and Chaisson	1997	-
						13.80	Lourens	et al.	2004	13.87	Turco	et al.	2002	14.60

M7	M7	T	Clavatorella	bermudezi	b	13.82	Lourens	et al.	2004	14.20	Pearson	and	Chaisson	1997	-
		T	Globorotalia	archeomenardii		13.87	Lourens	et al.	2004	13.87	Turco	et al.	2002	-	
		B	Fohsella	peripheroacuta		14.24	Lourens	et al.	2004	14.02	Turco	et al.	2002	14.80	
		B	Globorotalia	praemenardii		14.38	Lourens	et al.	2004	14.90	Pearson	and	Chaisson	1997	-
M6	M6	T	Praeorbulina	sicana	f	14.53	Lourens	et al.	2004	14.40	Shackleton	et al.	Chaisson	1999	14.80
		T	Globigerinatella	insueta		14.66	Lourens	et al.	2004	15.00	Pearson	and	Chaisson	1997	-
		B	Orbulina	suturalis		15.10	This study			15.10	Berggren	et al.	1995 b	15.10	
		B	Clavatorella	bermudezi		15.73	This study			15.80	Pearson	and	Chaisson	1997	-
M5b	M5b	B	Praeorbulina	circularis	e	15.96	This study			16.00	Berggren	et al.	1995 b	16.00	
		B	Globorotalia	archeomenardii		16.26	Lourens	et al.	2004	16.20	Pearson	and	Chaisson	1997	-
		B	Praeorbulina	glomerosa		16.27	Lourens	et al.	2004	16.20	Shackleton	et al.	1999	16.10	
		B	Praeorbulina	curva		16.28	This study			16.30	Berggren	et al.	1995 b	16.30	
M5a	M5a	B	Praeorbulina	sicana	sensu stricto	16.38	This study			16.40	Berggren	et al.	1995 b	16.40	
M4b	M4b	B	Fohsella	birnageae		16.69	This study			16.70	Berggren	et al.	1995 b	16.70	
M4a	M4a	B	Globorotalia	zealandica		h	17.26	This study			17.30	Berggren	et al.	1995 b	16.30
		T	Catapsydrax	dissimilis			17.54	Lourens	et al.	2004	17.50	Shackleton	et al.	1999	17.30
		B	Globigerinatella	insueta s. str.	17.59		Lourens	et al.	2004	17.40	Pearson	and	Chaisson	1997	(18.8)
		B	Globorotalia	praescitula	18.26		This study			18.50	Berggren	et al.	1995 b	18.50	
	M3	T	Globoquadrina	binaiensis		19.09	Lourens	et al.	2004	19.10	Pearson	and	Chaisson	1997	-
		B	Globigerinatella	sp.		19.30	Lourens	et al.	2004	20.20	Pearson	and	Chaisson	1997	-
		B	Globigerinoides	altiaperturus		20.03	This study			20.50	Berggren	et al.	1995 b	20.50	
		T	Tenuitella	munda		20.78	This study			21.40	Berggren	et al.	1995	21.40	
M2	M2	T	Globigerina	angulisuturalis		20.94	This study			21.60	Berggren	et al.	1995 b	21.60	
		T	'Paragloborotalia'	kugleri		21.12	Lourens	et al.	2004	21.00	Shackleton	et al.	1999	21.50	
M1b	M1b	T	'Paragloborotalia'	pseudokugleri		21.31	Lourens	et al.	2004	21.10	Shackleton	et al.	1999	21.60	
		B	Globoquadrina	dehiscens		22.44	This study			23.20	Berggren	et al.	1995 b	23.20	
M1a	M1a	T	Globigerina	ciperoensis	sensu lato	22.90	Lourens	et al.	2004	22.80	Shackleton	et al.	1999	-	
		B	Globigerinoides	trilobus		22.96	Lourens	et al.	2004	22.90	Shackleton	et al.	1999	-	
		B	'Paragloborotalia'	kugleri		22.96	Lourens	et al.	2004	22.90	Shackleton	et al.	1999	23.80	
		Oligocene/Miocene boundary						23.03	Lourens	et al.	2004	23.80	Berggren	et al.	1995 b

Notes

- a Calibrated to Lourens et al. (2004)
- b Calibration of Turco et al. (2002) on Astronomical Time Scale of Shackleton and Crowhurst (1997)
- c Replaces the and/or non zone of (1) BKSA95
- e Note mistake in Lourens et al. (2004) Table A2.3, where Bottom *Praeorbulina circularis* should read Top *Praeorbulina circularis*
- f Note mistake in Shackleton et al. (1999) where 14.8 Ma should read 15.8 Ma. This mistake is also propagated in Lourens et al. (2004)
- g Traditionally used as base N22, this datum is considered highly diachronous between ocean basins (see Dowsett, 1988).
- h Note mistake in BKSA95 (page 174) where LAD should read FAD
- i Note mistake in Berggren et al. (1995a, Table 6) where 1.6 Ma should read 1.77 Ma
- X Change in coiling direction

Wade et al Table 4

Zone (BKSA95 & BP05) (base)	Zone (This study) (base)	Datum		Age (Ma)		Astronomical		Published		Reference	Berggren et al.		Berggren & Pearson (2005)
				This Study ^a	Age ^b	Age ^b	Age ^b	Age (Ma)	Age (Ma)		(1995a,b)	(1995a,b)	
		T	Tenuitella	gemma	23.6	23.5		24.3	Berggren	et al.	1995 b	24.3	-
		LCO	Globigerinoides	primordius	23.6	23.5		24.3	Berggren	et al.	1995 b	24.3	-
	O7	B	'Paragloborotalia'	pseudokugleri	25.4	25.2		25.9	Berggren	et al.	1995 b	25.9	-
		B	Globigerinoides	primordius	26.3	26.1		26.7	Berggren	et al.	1995 b	26.7	-
O6	O6	T	Paragloborotalia	opima	27.3	26.9		27.5	Wade	et al.	2007	27.1	27.1
O5	O5	HCO	Chiloguembelina	cubensis	28.3	28.0		28.4	Wade	et al.	2007	28.5	28.5
O4	O4	B	Globigerina	angulisuturalis	29.5	29.2		29.4	Berggren	et al.	1995 b	29.4	29.4
		T	Subbotina	angiporoides	30.1	29.8		30.0	Berggren	et al.	1995 b	30.0	30.0
O3	O3	T	Turborotalia	ampliapertura	30.4	30.3		30.3	Berggren	et al.	1995 b	30.3	30.3
		B	Paragloborotalia	opima	30.8	30.8		30.6	Berggren	et al.	1995 b	30.6	-
O2	O2	T	Pseudohastigerina	nagewichiensis	32.2	32.0		32.0	Berggren	et al.	1995 b	32.0	32.0
<i>Eocene/Oligocene boundary</i>										<i>33.7 Berggren et al. 1995 b</i>	-	-	-
O1	O1	T	Hantkenina	alabamensis	33.9	33.8		33.7	Berggren	and Pearson	2005	33.7	33.7
		HCO	Pseudohastigerina	micra	33.9	33.8		33.7	Wade	and Pearson	2008	-	-
		T	Turborotalia	cerroazulensis	34.0	33.9		33.8	Berggren	and Pearson	2005	33.8	33.8
E16	E16	T	Globigerinatheka	index	34.5	34.5		34.3	Berggren	and Pearson	2005	34.3	34.3
		B	Turborotalia	cunialensis	35.4	35.4		35.3	Berggren	and Pearson	2005	35.2	35.3
E15	E15	T	Globigerinatheka	semiinvoluta	35.8	35.8		35.8	Berggren	and Pearson	2005	35.3	35.8
		B	Globigerinatheka	semiinvoluta	37.7	38.0		38.0	Wade		2004	38.4	-
E14	E14	T	Morozovelloides	crassatus	37.7	38.1		38.0	Wade		2004	38.1	38.0
		T	Acarinina	mcgowrani	37.7	38.1		38.0	Wade		2004	-	-
		T	Turborotalia	frontosa	38.8	39.3		39.3	Berggren	et al.	1995 b	39.3	-
E13	E13	T	Orbulinoides	beckmanni	39.4	40.0		40.0	Wade		2004	40.1	40.0
E12	E12	B	Orbulinoides	beckmanni	39.8	40.8		40.5	Berggren	et al.	1995 b	40.5	40.5
		T	Acarinina	bullbrookii	39.8	40.8		40.5	Berggren	et al.	1995 b	40.5	-
E11	E11	T	Guembilitrioides	nuttalli	(41.4)	-		(42.3)	Berggren	and Pearson	2005	-	(42.3)
		B	Turborotalia	pomeroli	41.5	-		42.4	Berggren	et al.	1995 b	42.4	-
		B	Globigerinatheka	index	41.9	-		42.9	Berggren	et al.	1995 b	42.9	-
		B	Morozovelloides	lehneri	42.5	-		43.5	Berggren	et al.	1995 b	43.5	-
E10	E10	T	Morozovella	aragonensis	42.6	-		43.6	Berggren	et al.	1995 b	43.6	43.6
E9	E9	B	Globigerinatheka	kugleri	(43.4)	-		-	Pearson	et al.	2004	45.8	45.8
		B	Hantkenina	singanoae	43.5	-		43.5	Payros	et al.	2009	49.0	-
E8	E8	B	Guembilitrioides	nuttalli	45.5	-		45.5	Payros	et al.	2009	-	49.0
	E7b	B	Turborotalia	frontosa	48.6	-		48.6	Payros	et al.	2009	-	-
E7	E7a	B	Acarinina	cuneicamerata	50.3	-		50.4	Hancock	et al.	2002	-	50.4
		B	Planorotalites	palmerae	50.3	-		50.4	Berggren	et al.	1995 b	50.4	-


E6	E6	T	Morozovella	subbotinae	50.8	-	50.8	Berggren	and	Pearson	2005	-	50.8
E5	E5	B	Morozovella	aragonensis	52.3	-	52.3	Berggren	et	al.	1995 b	52.3	52.3
		T	Morozovella	marginodentata	52.8	-	52.5	Berggren	et	al.	1995 b	52.5	-
		T	Morozovella	lensiformis	53.1	-	52.7	Berggren	et	al.	1995 b	52.7	-
		T	Morozovella	aequa	54.0	-	53.6	Berggren	et	al.	1995 b	53.6	-
E4	E4	B	Morozovella	formosa	54.4	-	54.0	Berggren	et	al.	1995 b	54.0	54.0
		B	Morozovella	lensiformis	54.4	-	54.0	Berggren	et	al.	1995 b	54.0	-
E3	E3	T	Morozovella	velascoensis	54.9	-	54.5	Berggren	and	Pearson	2005	54.7	54.5
		T	Morozovella	acuta	55.1	-	54.7	Berggren	et	al.	1995 b	54.7	-
		B	Morozovella	gracilis	55.1	-	54.7	Berggren	et	al.	1995 b	54.7	-
		B	Igorina	broedermanni	55.1	-	54.7	Berggren	et	al.	1995 b	54.7	-
		B	Morozovella	marginodentata	55.2	-	54.8	Berggren	et	al.	1995 b	54.8	-
E2	E2	B	Pseudohastigerina	wilcoxensis	55.7	-	55.4	Berggren	and	Pearson	2005	-	55.4
		B	Globanomalina	australiformis	55.8	-	55.5	Berggren	et	al.	1995 b	55.5	-
E1	E1	B	Acarinina	sibaiyaensis	55.8	-	55.5	Berggren	and	Pearson	2005	-	55.5
					55.8	-	55.5	Ouda	and	Aubry	2003		
P5	P5	T	Globanomalina	pseudomenardii	56.7	-	55.9	Berggren	et	al.	1995 b	55.9	55.9
		B	Morozovella	subbotinae	56.7	-	55.9	Berggren	et	al.	1995 b	55.9	-
		T	Acarinina	mckannai	57.1	-	56.3	Berggren	et	al.	1995 b	56.3	-
		T	Acarinina	acarinata	57.1	-	56.3	Berggren	et	al.	1995 b	56.3	-
P4c	P4c	B	Acarinina	soldadoensis	57.3	-	56.5	Berggren	et	al.	1995 b	56.5	56.5
		B	Acarinina	coalingensis	57.3	-	56.5	Berggren	et	al.	1995 b	56.5	-
		B	Morozovella	aequa	57.3	-	56.5	Berggren	et	al.	1995 b	56.5	-
		T	Acarinina	subsphaerica	57.9	-	57.1	Berggren	et	al.	1995 b	57.1	57.1
		B	Acarinina	mckannai	59.9	-	59.1	Berggren	et	al.	1995 b	59.1	-
P4b	P4b	T	Parasubbotina	variospira	60.0	-	59.2	Berggren	et	al.	1995 b	59.2	59.2
		B	Acarinina	acarinata	60.0	-	59.2	Berggren	et	al.	1995 b	59.2	-
		B	Acarinina	subsphaerica	60.0	-	59.2	Berggren	et	al.	1995 b	59.2	-
P4a	P4a	B	Globanomalina	pseudomenardii	60.2	-	59.4	Berggren	et	al.	2000	59.2	59.4
P3b	P3b	B	Igorina	albeari	60.8	-	60.0	Berggren	et	al.	1995 b	60.0	60.0
		B	Morozovella	velascoensis	60.8	-	60.0	Berggren	et	al.	1995 b	60.0	-
		B	Morozovella	conicotruncata	61.7	-	60.9	Berggren	et	al.	1995 b	60.9	-
P3a	P3a	B	Morozovella	angulata	61.7	-	61.0	Berggren	et	al.	1995 b	61.0	61.0
		B	Igorina	pusilla	61.7	-	61.0	Berggren	et	al.	1995 b	61.0	-
		B	Morozovella	praeangulata	61.9	-	61.2	Berggren	et	al.	1995 b	61.2	-
		B	Globanomalina	imitata	(62.0)	-	(61.3)	Berggren	et	al.	1995 b	(61.3)	-
P2	P2	B	Praemurica	uncinata	62.1	-	61.4	Berggren	and	Pearson	2005	61.2	61.4
P1c	P1c	B	Globanomalina	compressa	63.5	-	62.9	Berggren	and	Pearson	2005	63.0	62.9
		B	Praemurica	inconstans	63.5	-	62.9	Berggren	and	Pearson	2005	63.0	62.9
		B	Parasubbotina	variata	63.6	-	63.0	Berggren	et	al.	1995 b	63.0	-

P1b	P1b	B	Subbotina triloculinoides	64.7	-	64.3 Berggren et al. 1995 b	64.3	64.3
P1a	P1a	T	Parvularugoglobigerina eugubina	65.2	-	64.8 Berggren and Pearson 2005	64.7	64.8
		B	Parvularugoglobigerina extensa	65.4	-	64.9 Olsson et al. 1999	-	-
Pα	Pα	B	Parvularugoglobigerina eugubina	65.46	-	64.97 Berggren et al. 1995 b	64.97	64.97
P0	P0	T	Globotruncana	65.5	-	65.0 Berggren et al. 1995 b	65.0	65.0


Notes

a	Calibrated to Gradstein et al. (2004)
b	Calibrated to Pälike et al. (2006)
c	Replaces the and/or non zone of Berggren & Pearson (2005)
d	Note mistake in Berggren & Pearson (2005)
HCO	Highest common occurrence
LCO	Lowest common occurrence
(43.4)	Estimated age

Wade et al.
Figure 1




Age (Ma; Cande and Kent, 1995)




Wade et al. Figure 2a


Age (Ma; Cande and Kent, 1995)



Age (Ma; Cande and Kent, 1995)




Age (Ma; astronomical)





Wade et al. Fig. 3a

Age (Ma; astronomical)




Wade et al. Fig 3b






Age (Ma; Gradstein et al., 2004)



Stage	Blow (1969)	Berggren et al. (1995)	Wade et al. (this study)	Datum Events
middle Miocene	N14	M11	M11	
	N13	M10	M10	
	N12	M9b	M9b	
		M8/M9a	M9a	
	N11	M7	M8	
	N10		M7	
	N9		M6	
	N8	M5	M5	

Wade et al.
Figure 5



Wade et al. Figure 6